Noname manuscript No.
(will be inserted by the editor)

arXiv:1005.0336v3 [math.CA] 21 Jun 2010

Zeros of Orthogonal Polynomials Generated by Canonical Perturbations on Standard Measure

Edmundo J. Huertas · **Francisco Marcellán** ·
Fernando R. Rafaeli.

Received: date / Accepted: date

Abstract In this contribution, the behavior of zeros of orthogonal polynomials associated with canonical linear spectral transformations of measures supported in the real line is analyzed. An electrostatic interpretation of them is given.

Keywords Orthogonal polynomials · Connection formula · distribution of zeros · interlacing · monotonicity · asymptotic behavior · electrostatics interpretation

Mathematics Subject Classification (2000) Primary 30C15

1 Introduction

1.1 Basic theory of orthogonal polynomials on the real line

Let μ be a positive Borel measure supported on a subset Σ of the real line with infinitely many points and such that

$$\int_{\Sigma} |x|^n d\mu(x) < \infty,$$

This work has been done in the framework of a joint project of a joint project of Dirección General de Investigación, Ministerio de Educación y Ciencia of Spain and the Brazilian Science Foundation CAPES, Project CAPES/DGU 160/08. The work of the first and second authors has been supported by Dirección General de Investigación, Ministerio de Ciencia e Innovación of Spain, grant MTM2009-12740-C03-01. The work of the third author has been supported by FAPESP under grant 2009/14776-5.

Edmundo J. Huertas
Departamento de Matemáticas, Escuela Politécnica Superior
Universidad Carlos III, Leganés-Madrid, Spain
E-mail: ehuertas@math.uc3m.es

Francisco Marcellán
Departamento de Matemáticas, Escuela Politécnica Superior
Universidad Carlos III, Leganés-Madrid, Spain
E-mail: pacomarc@ing.uc3m.es second address

Fernando R. Rafaeli
Departamento de Ciências de Computação e Estatística
IBILCE, Universidade Estadual Paulista, Brazil
E-mail: fe_ro_rafaeli@yahoo.com.br

for every $n \in \mathbb{Z}_+$.

Given a measure μ, we define the standard inner product $\langle \cdot, \cdot \rangle_\mu : \mathbb{P} \times \mathbb{P} \to \mathbb{R}$ by

$$\langle p, q \rangle_\mu = \int_\Sigma p(x)q(x)d\mu(x), \quad p, q \in \mathbb{P}, \tag{1}$$

where $\mathbb{P}$ is the linear space of the polynomials with real coefficients, and the corresponding norm $\| \cdot \|_\mu : \mathbb{P} \to [0, \infty)$ is given, as usual, by

$$\|p\|_\mu = \sqrt{\int_\Sigma |p(x)|^2 d\mu(x)}, \quad p \in \mathbb{P}. \tag{2}$$

Definition 1 *Let* $\{p_n(x)\}_{n\geq 0}$ *be a sequence of polynomials such that*

1. $p_n(x)$ *is a polynomial with degree exactly* n.
2. $\langle p_n, p_m \rangle_\mu = 0$, *for* $m \neq n$.

$\{p_n(x)\}_{n\geq 0}$ *is said to be a* ***sequence of standard orthogonal polynomials.*** *If the leading coefficient of* p_n *is* 1, *then the sequence is said to be a standard* ***monic orthogonal polynomial sequence (MOPS, in short).***

Proposition 1 *Each positive Borel measure* μ *determines uniquely a standard MOPS.*

A MOPS is generated by a three term recurrence relation. It will play an important role in the sequel.

Proposition 2 (Three Term Recurrence Relation TTRR) *Let* $\{p_n(x)\}_{n\geq 0}$ *be a MOPS. They satisfy a three-term recurrence relation*

$$p_{n+1}(x) = (x - \beta_n)p_n(x) - \gamma_n p_{n-1}(x), \quad n \geq 0, \tag{3}$$

with initial conditions $p_0(x) = 1$ *and* $p_{-1}(x) = 0$. *The recurrence coefficients are given by*

$$\beta_n = \frac{\langle xp_n, p_n \rangle_\mu}{\|p_n\|_\mu^2}, \ n \geq 0,$$

and

$$\gamma_n = \frac{\|p_n\|_\mu^2}{\|p_{n-1}\|_\mu^2} > 0, n \geq 1.$$

Next, we will introduce the *nth* Kernel associated with the MOPS $\{p_n(x)\}_{n\geq 0}$. It satisfies a reproducing property for every polynomial of degree at most n as well it can be represented in a simple way in terms of the polynomials $p_n(x)$ and $p_{n+1}(x)$ throught the Christoffel-Darboux Formula, that can be deduced in a straightforward way from the three term recurrence relation (see [3]).

Proposition 3 *Let* $\{p_n(x)\}_{n\geq 0}$ *be a MOPS. If we denote the nth Kernel polynomial by*

$$K_n(x, y) = \sum_{j=0}^{n} \frac{p_j(y)p_j(x)}{\|p_j\|_\mu^2}, \tag{4}$$

then, for every $n \in \mathbb{N}$,

$$K_n(x, y) = \frac{1}{\|p_n\|_\mu^2} \frac{p_{n+1}(x)p_n(y) - p_n(x)p_{n+1}(y)}{x - y}.$$

and, if $x = y$ we have the so called confluent form

$$K_n(x,x) = \frac{p'_{n+1}(x)p_n(x) - p'_n(x)p_{n+1}(x)}{\|p_n\|_\mu^2}.$$

The zeros of standard orthogonal polynomials are the nodes of the Gaussian quadrature rules and they have nice analytic properties. Next, we summarize some of them. For more information, see [3], [19], and [21]

Proposition 4 *Let μ be a positive Borel measure defined as above and $\{p_n(x)\}_{n\geq 0}$ the corresponding MOPS. Then,*

1. *For each $n \geq 1$, the polynomial $p_n(x)$ has n real and simple zeros in the interior of $C_0(\Sigma)$, the convex hull of Σ.*
2. *The zeros of $p_{n+1}(x)$ interlace with the zeros of $p_n(x)$.*
3. *Between any two consecutive zeros of $p_n(x)$ there is at least one zero of $p_m(x)$, for $m > n \geq 2$.*
4. *If $(\alpha, \beta) \subset C_0(\Sigma)$ with $(\alpha, \beta) \cap \Sigma = \emptyset$ then at most one zero of each polynomial $p_n(x)$ belongs to (α, β).*
5. *Each point of Σ attracts zeros of the MOPS. In other words, the zeros of a MOPS are dense in Σ.*

Next, we will analyze the behavior of zeros of polynomial of the form $f(x) = h_n(x) + cg_n(x)$

We need the following lemma concerning the behavior and the asymptotics of the zeros of linear combinations of two polynomials with interlacing zeros (see [2, Lemma 1] and [5, Lemma 3]).

Lemma 1 *Let $h_n(x) = a(x - x_1)\cdots(x - x_n)$ and $g_n(x) = b(x - \zeta_1)\cdots(x - \zeta_n)$ be polynomials with real and simple zeros, where a and b are real positive constants.*

(i) If

$$\zeta_1 < x_1 < \cdots < \zeta_n < x_n,$$

then, for any real constant $c > 0$, the polynomial

$$f(x) = h_n(x) + cg_n(x)$$

has n real zeros $\eta_1 < \cdots < \eta_n$ which interlace with the zeros of $h_n(x)$ and $g_n(x)$ in the following way

$$\zeta_1 < \eta_1 < x_1 < \cdots < \zeta_n < \eta_n < x_n.$$

Moreover, each $\eta_k = \eta_k(c)$ is a decreasing function of c and, for each $k = 1, \ldots, n$,

$$\lim_{c\to\infty} \eta_k = \zeta_k \quad \text{and} \quad \lim_{c\to\infty} c[\eta_k - \zeta_k] = \frac{-h_n(\zeta_k)}{g'_n(\zeta_k)}. \tag{5}$$

(ii) If

$$x_1 < \zeta_1 < \cdots < x_n < \zeta_n,$$

then, for any positive real constant $c > 0$, the polynomial

$$f(x) = h_n(x) + cg_n(x)$$

has n real zeros $\eta_1 < \cdots < \eta_n$ which interlace with the zeros of $h_n(x)$ and $g_n(x)$ as follows

$$x_1 < \eta_1 < \zeta_1 < \cdots < x_n < \eta_n < \zeta_n.$$

Moreover, each $\eta_k = \eta_k(c)$ is an increasing function of c and, for each $k = 1, \ldots, n$,

$$\lim_{c\to\infty} \eta_k = \zeta_k \quad \textit{and} \quad \lim_{c\to\infty} c[\zeta_k - \eta_k] = \frac{h_n(\zeta_k)}{g_n'(\zeta_k)}. \tag{6}$$

In the last years some attention has been paid to the so called canonical spectral transformations of measures. Some authors have analyzed them from the point of view of Stieltjes functions associated with such a kind of perturbations (see [23])or from the relation between the corresponding Jacobi matrices (see [24]). Our present contribution is focused on the behaviour of zeros of MOPS associated with such transformations of the measures. In particular, we are interested in the Christoffel and Uvarov transformations which are given as a multiplication of the measure by a positive linear polynomial in its support and the addition of a Dirac mass at a point outside the support, respectively.

The structure of the manuscript is as follows. In Section 2 the representation of the perturbed MOPS in terms of the initial ones is done. In Section 3 we analyze the behaviour of the zeros of the MOPS when an Uvarov transform is introduced. In particular, we obtain such a behavior when the mass N tends to infinity as well as we characterize the values of the mass N such the smallest (respectively, the largest) zero of these MOPS is located outside the support of the measure. In Section 4, we check these results in the cases of the Jacobi-type and Laguerre-type orthogogonal polynomials introduced by T. H. Koornwinder ([14]). Section 5 is devoted to the electrostatic interpretation of the zero distribution as equilibrium points in a logarithmic potential interaction under the action of an external field.We analyze such an equilibrium problem when the mass point is located on the boundary or in the exterior of the support of the measure, respectively.

2 Canonical perturbations of a measure

Let $\{p_n(x)\}_{n\geq 0}$ be the MOPS with respect to a positive Borel measure μ defined as above. We will consider some canonical perturbations of the measure, which are called spectral linear transformations (see [23] and [24]).

2.1 Christoffel perturbation

Let $\{p_n^*(a;x)\}_{n\geq 0}$ be the MOPS associated with the measure

$$d\mu^* = (x-a)d\mu,$$

with $a \notin C_0(\Sigma)$. This means that $p_n(a) \neq 0$ for every $n \geq 1$.

The polynomial $p_n^*(a;x)$ is given by (see [3, (7.3)])

$$p_n^*(a;x) = \frac{1}{x-a}\left[p_{n+1}(x) - \frac{p_{n+1}(a)}{p_n(a)}p_n(x)\right] = \frac{\|p_n\|_\mu^2}{p_n(a)}K_n(a,x), \tag{7}$$

i.e., $p_n^*(a;x)$ is a monic kernel polynomial. Notice that $p_n^*(a;a) \neq 0$. Let $x_{n,k}$ and $x_{n,k}^* := x_{n,k}^*(a)$ be the zeros of $p_n(x)$ and $p_n^*(a;x)$, respectively, all arranged in an increasing order, and assume that $C_0(\Sigma) = [\xi, \eta]$. From (7) and Proposition 4 (item 2) the following interlacing property of these zeros holds

(i) If $a \leq \xi$, then $p_{n+1}(a)/p_n(a) < 0$, and

$$Sign\left[p_n^*(a;x_{n+1,k})\right] = Sign\left[p_n(x_{n+1,k})\right], \; k = 1, \dots, n+1,$$

as well as

$$Sign\left[p_n^*(a;x_{n,k})\right] = Sign\left[p_{n+1}(x_{n,k})\right], \; k = 1, \dots, n.$$

Therefore

$$x_{n+1,1} < x_{n,1} < x_{n,1}^* < x_{n+1,2} < \cdots < x_{n,n} < x_{n,n}^* < x_{n+1,n+1}; \tag{8}$$

(ii) If $a \geq \eta$, then $p_{n+1}(a)/p_n(a) > 0$ and, as a consequence,

$$Sign\left[p_n^*(a;x_{n+1,k})\right] = Sign\left[p_n(x_{n+1,k})\right], \; k = 1, \dots, n+1,$$

and

$$Sign\left[p_n^*(a;x_{n,k})\right] = -Sign\left[p_{n+1}(x_{n,k})\right], \; k = 1, \dots, n.$$

Therefore

$$x_{n+1,1} < x_{n,1}^* < x_{n,1} < \cdots < x_{n+1,n} < x_{n,n}^* < x_{n,n} < x_{n+1,n+1}. \tag{9}$$

2.2 Iterated-Christoffel

Let $\{p_n^{**}(a;x)\}_{n\geq 0}$ be the MOPS associated with the measure

$$d\mu^{**} = (x-a)^2 d\mu,$$

with $a \notin C_0(\Sigma)$. Using (7) we deduce that

$$\begin{aligned} p_n^{**}(a;x) &= \frac{1}{x-a}\left[p_{n+1}^*(a;x) - \frac{p_{n+1}^*(a;a)}{p_n^*(a;a)} p_n^*(a;x)\right] \\ &= \frac{\|p_n^*\|_{\mu^*}^2}{p_n^*(a;a)} K_n^*(a,x) \\ &= \frac{1}{(x-a)^2}\left[p_{n+2}(x) - d_n p_{n+1}(x) + e_n p_n(x)\right], \end{aligned} \tag{10}$$

where $K_n^*(a,x)$ is the Kernel polynomial associated with $p_n^*(a;x)$, and

$$\begin{aligned} d_n &= \frac{p_{n+2}(a)}{p_{n+1}(a)} + \frac{p_{n+1}^*(a;a)}{p_n^*(a;a)} = \frac{p_{n+2}(a) + p_n(a)}{p_{n+1}(a)} e_n, \\ e_n &= \frac{p_{n+1}^*(a;a)}{p_n^*(a;a)} \frac{p_{n+1}(a)}{p_n(a)} = \frac{\|p_{n+1}\|_\mu^2}{\|p_n\|_\mu^2} \frac{K_{n+1}(a,a)}{K_n(a,a)} > 0. \end{aligned}$$

Notice that $p_n^{**}(a;a) \neq 0$. Let denote by $x_{n,k}^{**} := x_{n,k}^{**}(a)$ the zeros of $p_n^{**}(a;x)$, arranged in an increasing order. Then replacing (3) in (10) we obtain

$$p_n^{**}(a;x) = \frac{1}{(x-a)^2}\left[(x-\beta_{n+1}-d_n)p_{n+1}(x) + (e_n - \gamma_{n+1})p_n(x)\right]. \tag{11}$$

On the other hand,

$$e_n - \gamma_{n+1} = \frac{\|p_{n+1}\|_\mu^2}{\|p_n\|_\mu^2}\left(\frac{K_{n+1}(a,a)}{K_n(a,a)} - 1\right) > 0. \tag{12}$$

Evaluating $p_n^{**}(a;x)$ at the zeros $x_{n+1,k}$, from (11) and (12), we get

$$Sign\left[p_n^{**}(a;x_{n+1,k})\right] = Sign\left[p_n(x_{n+1,k})\right],\ k = 1,\ldots,n+1. \tag{13}$$

Thus, from (13) and Proposition 4 (item 2) we obtain the following interlacing property:

Theorem 1 *The inequalities*

$$x_{n+1,1} < x_{n,1}^{**} < x_{n+1,2} < x_{n,2}^{**} < \cdots < x_{n+1,n} < x_{n,n}^{**} < x_{n+1,n+1} \tag{14}$$

hold for every $n \in \mathbb{N}$.

2.3 Uvarov perturbation

Let $\{p_n^N(a;x)\}_{n\geq 0}$ be the MOPS associated with the measure

$$d\mu_N = d\mu + N\delta_a,$$

with $N \in \mathbb{R}_+$, δ_a the Dirac delta function in $x = a$, and $a \notin C_0(\Sigma)$. R. Álvarez-Nodarse, F. Marcellán and J. Petronilho [1, (8)] obtained the following representation for such polynomials in terms of the MOPS $\{p_n(x)\}_{n\geq 0}$ (see also [22].)

$$p_n^N(a;x) = p_n(x) - \frac{Np_n(a)}{1 + NK_{n-1}(a,a)}K_{n-1}(a,x). \tag{15}$$

Next, we give another connection formula for the Uvarov's orthogonal polynomials $p_n^N(a;x)$ using the standard orthogonal polynomials $p_n(x)$ and the Iterated-Christoffel's orthogonal polynomials $p_n^{**}(a;x)$.

Theorem 2 (Connection Formula) *The polynomials* $\{\widehat{p}_n^N(a;x)\}_{n\geq 0}$, *with* $\widehat{p}_n^N(a;x) = k_n p_n^N(a;x)$, *can be represented as*

$$\widehat{p}_n^N(a;x) = p_n(x) + NB_n(x-a)p_{n-1}^{**}(a;x), \tag{16}$$

with

$$B_n = \frac{-p_n(a)}{\langle x-a, p_{n-1}^{**}\rangle_\mu} = K_{n-1}(a,a) > 0 \tag{17}$$

and $k_n = 1 + NB_n$.

Proof In order to prove the orthogonality of the polynomials defined by (16), we deal with the basis $1,(x-a),(x-a)^2,\ldots,(x-a)^n$ of the linear space of polynomials of degree at most n. Then,

$$\begin{aligned}
\langle 1,\widehat{p}_n^N\rangle_{\mu_N} &= \langle 1,p_n\rangle_\mu + NB_n\langle 1,(x-a)p_{n-1}^{**}\rangle_\mu + Np_n(a)=0\\
\langle (x-a),\widehat{p}_n^N\rangle_{\mu_N} &= \langle (x-a),p_n\rangle_\mu + NB_n\langle 1,p_{n-1}^{**}\rangle_{\mu^{**}}=0\\
&\vdots\\
\langle (x-a)^{n-1},\widehat{p}_n^N\rangle_{\mu_N} &= \langle (x-a)^{n-1},p_n\rangle_\mu + NB_n\langle (x-a)^{n-2},p_{n-1}^{**}\rangle_{\mu^{**}}=0,
\end{aligned}$$

and, finally,

$$\begin{aligned}
\langle (x-a)^{n},\widehat{p}_n^N\rangle_{\mu_N} &= \langle (x-a)^{n},p_n\rangle_\mu + NB_n\langle (x-a)^{n-1},p_{n-1}^{**}\rangle_{\mu^{**}}>0\\
&= \|p_n\|_\mu^2 + NB_n\|p_{n-1}^{**}\|_{\mu^{**}}^2>0.
\end{aligned}$$

In order to prove (17), from (10) and (7) we get

$$\begin{aligned}
\langle x-a,p_{n-1}^{**}\rangle_\mu &= \int (x-a)\,p_{n-1}^{**}(a;x)d\mu(x)\\
&= \int (x-a)\,\frac{1}{x-a}\left[p_n^*(a;x)-\frac{p_n^*(a;a)}{p_{n-1}^*(a;a)}p_{n-1}^*(a;x)\right]d\mu(x)\\
&= \int p_n^*(a;x)d\mu(x)-\frac{p_n^*(a;a)}{p_{n-1}^*(a;a)}\int p_{n-1}^*(a;x)d\mu(x)\\
&= \frac{\|p_n\|_\mu^2}{p_n(a)}\int K_n(a;x)d\mu(x)\\
&\quad -\frac{\|p_n\|_\mu^2}{p_n(a)}\frac{K_n(a;a)}{K_{n-1}(a;a)}\frac{p_{n-1}(a)}{\|p_{n-1}\|_\mu^2}\frac{\|p_{n-1}\|_\mu^2}{p_{n-1}(a)}\int K_{n-1}(a;x)d\mu(x)\\
&= \frac{\|p_n\|_\mu^2}{p_n(a)}-\frac{\|p_n\|_\mu^2}{p_n(a)}\frac{K_n(a;a)}{K_{n-1}(a;a)}\\
&= \frac{\|p_n\|_\mu^2}{p_n(a)}\left(1-\frac{K_n(a;a)}{K_{n-1}(a;a)}\right).
\end{aligned}$$

Thus

$$B_n=\frac{-p_n(a)}{\langle x-a,p_{n-1}^{**}\rangle_\mu}=\frac{p_n^2(a)}{\|p_n\|_\mu^2\left[K_n(a,a)/K_{n-1}(a,a)-1\right]}=K_{n-1}(a,a)>0$$

3 The Zeros

We call the attention of the reader on the fact that the constant B_n defined as above does not depend on N. For this reason, the connection formula (16) is very useful in order to obtain results about monotonicity, asymptotics, and speed of convergence for the zeros of $p_n^N(a;x)$ in terms of the mass N. Indeed, let assume that $x_{n,k}^N := x_{n,k}^N(a), k=1,2,...,n$, are the zeros of $p_n^N(a;x)$. Thus, from (14), (16), and Lemma 1, we immediately conclude that

Theorem 3 *If $C_0(\Sigma)=[\xi,\eta]$ and $a\le\xi$, then*

$$a<x_{n,1}^N<x_{n,1}<x_{n-1,1}^{**}<x_{n,2}^N<x_{n,2}<\cdots<x_{n-1,n-1}^{**}<x_{n,n}^N<x_{n,n}. \tag{18}$$

Moreover, each $x_{n,k}^N$ is a decreasing function of N and, for each $k=1,\dots,n-1$,

$$\lim_{N\to\infty}x_{n,1}^N=a,\quad \lim_{N\to\infty}x_{n,k+1}^N=x_{n-1,k}^{**}, \tag{19}$$

as well as

$$\begin{aligned}\lim_{N\to\infty}N[x_{n,1}^N-a]&=\frac{-p_n(a)}{B_np_{n-1}^{**}(a;a)},\\ \lim_{N\to\infty}N[x_{n,k+1}^N-x_{n-1,k}^{**}]&=\frac{-p_n(x_{n-1,k}^{**})}{B_n(x_{n-1,k}^{**}-a)[p_{n-1}^{**}(a;x)]'_{x=x_{n-1,k}^{**}}}.\end{aligned} \tag{20}$$

Theorem 4 *If $C_0(\Sigma)=[\xi,\eta]$ and $a\ge\eta$, then*

$$x_{n,1}<x_{n,1}^N<x_{n-1,1}^{**}<\cdots<x_{n,n-1}<x_{n,n-1}^N<x_{n-1,n-1}^{**}<x_{n,n}<x_{n,n}^N<a. \tag{21}$$

Moreover, each $x_{n,k}^N$ is an increasing function of N and, for each $k=1,\dots,n-1$,

$$\lim_{N\to\infty}x_{n,n}^N=a,\quad \lim_{N\to\infty}x_{n,k}^N=x_{n-1,k}^{**}, \tag{22}$$

and

$$\begin{aligned}\lim_{N\to\infty}N[a-x_{n,n}^N]&=\frac{p_n(a)}{B_np_{n-1}^{**}(a;a)},\\ \lim_{N\to\infty}N[x_{n-1,k}^{**}-x_{n,k}^N]&=\frac{p_n(x_{n-1,k}^{**})}{B_n(x_{n-1,k}^{**}-a)[p_{n-1}^{**}(a;x)]'_{x=x_{n-1,k}^{**}}}.\end{aligned} \tag{23}$$

Notice that the mass point a attracts one zero of $p_n^N(a;x)$, i.e. when $N\to\infty$, it captures either the smallest or the largest zero, according to the location of the point a with respect to the support of the measure μ.

3.1 The Minimum Mass

When either $a<\xi$ or $a>\eta$, at most one of the zeros of $p_n^N(a;x)$ is located outside of $C_0(\Sigma)=[\xi,\eta]$. In the next results, we will give explicitly the value N_0 of the mass such that for $N>N_0$ this situation occurs, i.e, one of the zeros is located outside $[\xi,\eta]$.

Corollary 1 *If $C_0(\Sigma)=[\xi,\eta]$ and $a<\xi$, then the smallest zero $x_{n,1}^N=x_{n,1}^N(a)$ satisfies*

$$\begin{aligned}x_{n,1}^N&>\xi,\quad \text{for}\quad N<N_0,\\ x_{n,1}^N&=\xi,\quad \text{for}\quad N=N_0,\\ x_{n,1}^N&<\xi,\quad \text{for}\quad N>N_0,\end{aligned}$$

where

$$N_0=N_0(n,a,\xi)=\frac{-p_n(\xi)}{K_{n-1}(a,a)\,(\xi-a)p_{n-1}^{**}(a;\xi)}>0.$$

Proof In order to deduce the location of $x_{n,1}^N$ with respect to the point $x = \xi$, it is enough to observe that $p_n^N(\xi) = 0$ if and only if $N = N_0$.

Corollary 2 *If $C_0(\Sigma) = [\xi, \eta]$ and $a > \eta$, then the largest zero $x_{n,n}^N = x_{n,n}^N(a)$ satisfies*

$$x_{n,n}^N < \eta, \quad \text{for} \quad N < N_0,$$

$$x_{n,n}^N = \eta, \quad \text{for} \quad N = N_0,$$

$$x_{n,n}^N > \eta, \quad \text{for} \quad N > N_0,$$

where

$$N_0 = N_0(n, a, \eta) = \frac{-p_n(\eta)}{K_{n-1}(a, a)(\eta - a)p_{n-1}^{**}(a; \eta)} > 0.$$

Proof In order to find the location of $x_{n,n}^N$ with respect to the point $x = \eta$, notice that $p_n^N(\eta) = 0$ if and only if $N = N_0$.

4 Application to classical measures

4.1 Jacobi type (Jacobi-Koornwinder) orthogonal polynomials

First, we will consider $p_n(x) = P_n^{\alpha,\beta}(x)$, the classical monic Jacobi polynomials, which are orthogonal with respect to the measure $d\mu_{\alpha,\beta} = (1-x)^\alpha(1+x)^\beta dx$, $\alpha, \beta > -1$, supported on $[-1, 1]$. We consider the Uvarov perturbations on $\mu_{\alpha,\beta}$ with either $a = -1$ or $a = 1$, and $M, N \geq 0$.

$$d\mu_M = d\mu_{\alpha,\beta} + M\delta_{-1}, \tag{24}$$

$$d\mu_N = d\mu_{\alpha,\beta} + N\delta_1. \tag{25}$$

Such orthogonal polynomials were first studied by T. H. Koornwinder (see [14]), in 1984. There, he adds simultaneously two Dirac delta functions at the end points $x = -1$ and $x = 1$, that is,

$$d\mu_{M,N} = d\mu_{\alpha,\beta} + M\delta_{-1} + N\delta_1.$$

Let $\{P_n^{\alpha,\beta,M}(x)\}_{n\geq 0}$ and $\{P_n^{\alpha,\beta,N}(x)\}_{n\geq 0}$ denote the sequences of orthogonal polynomials with respect (24) and (25), with the normalization pointed out in Theorem 2, respectively. Then, the connection formulas are

$$P_n^{\alpha,\beta,M}(x) = P_n^{\alpha,\beta}(x) + MK_{n-1}(-1,-1)(x+1)P_{n-1}^{\alpha,\beta+2}(x)$$

and

$$P_n^{\alpha,\beta,N}(x) = P_n^{\alpha,\beta}(x) + NK_{n-1}(1,1)(x-1)P_{n-1}^{\alpha+2,\beta}(x). \tag{26}$$

It is straightforward to see that

$$K_{n-1}(-1,-1) = \frac{1}{2^{\alpha+\beta+1}} \frac{\Gamma(n+\beta+1)\Gamma(n+\alpha+\beta+1)}{\Gamma(n)\Gamma(\beta+1)\Gamma(\beta+2)\Gamma(n+\alpha)}$$

and

$$K_{n-1}(1,1) = \frac{1}{2^{\alpha+\beta+1}} \frac{\Gamma(n+\alpha+1)\Gamma(n+\alpha+\beta+1)}{\Gamma(n)\Gamma(\alpha+1)\Gamma(\alpha+2)\Gamma(n+\beta)}.$$

Recently, several authors ([1], [5], [8]) have been contributed to the analysis of the behavior of the zeros of $P_n^{\alpha,\beta,M}(x)$, and $P_n^{\alpha,\beta,N}(x)$.

Let denote by $(x_{n,k}^M(\alpha))$, $(x_{n,k}^N(\alpha))$, and $(x_{n,k}(\alpha))$ the zeros of $P_n^{\alpha,\beta,M}(x)$, $P_n^{\alpha,\beta,N}(x)$, and $P_n^{\alpha,\beta}(x)$, respectively, all arranged in an increasing order. Then, applying the results of Section 3, we obtain

Theorem 5 *The inequalities*

$$-1 < x_{n,1}^M(\alpha,\beta) < x_{n,1}(\alpha,\beta) < x_{n-1,1}(\alpha,\beta+2) < x_{n,2}^N(\alpha,\beta) < x_{n,2}(\alpha,\beta) < \cdots$$
$$< x_{n-1,n-1}(\alpha,\beta+2) < x_{n,n}^N(\alpha,\beta) < x_{n,n}(\alpha,\beta)$$

hold for every $\alpha,\beta > -1$. *Moreover, each* $x_{n,k}^M(\alpha,\beta)$ *is a decreasing function of* M *and, for each* $k = 1,\ldots,n-1$,

$$\lim_{M\to\infty} x_{n,1}^M(\alpha,\beta) = -1, \quad \lim_{M\to\infty} x_{n,k+1}^M(\alpha,\beta) = x_{n-1,k}(\alpha,\beta+2), \tag{27}$$

and

$$\lim_{M\to\infty} M[x_{n,1}^M(\alpha,\beta)+1] = h_n(\alpha,\beta),$$

$$\lim_{M\to\infty} M[x_{n,k+1}^M(\alpha,\beta) - x_{n-1,k}(\alpha,\beta+2)] = \frac{\big[1 - x_{n-1,k}(\alpha,\beta+2)\big]\, h_n(\alpha,\beta)}{2(\beta+2)},$$

where

$$h_n(\alpha,\beta) = \frac{2^{\alpha+\beta+2}\Gamma(n)\Gamma(\beta+2)\Gamma(\beta+3)\Gamma(n+\alpha)}{\Gamma(n+\beta+2)\Gamma(n+\alpha+\beta+2)}.$$

Proof It remains to show the limits above. From (20)

$$\lim_{M\to\infty} M[x_{n,1}^M(\alpha,\beta)+1] = \frac{-P_n^{\alpha,\beta}(-1)}{K_{n-1}(-1,-1)P_{n-1}^{\alpha,\beta+2}(-1)}$$

Since

$$P_n^{\alpha,\beta}(-1) = \frac{(-1)^n 2^n \Gamma(n+\beta+1)\Gamma(n+\alpha+\beta+1)}{\Gamma(\beta+1)\Gamma(2n+\alpha+\beta+1)}.$$

and

$$K_{n-1}(-1,-1) = \frac{1}{2^{\alpha+\beta+1}} \frac{\Gamma(n+\beta+1)\Gamma(n+\alpha+\beta+1)}{\Gamma(n)\Gamma(\beta+1)\Gamma(\beta+2)\Gamma(n+\alpha)}$$

we obtain

$$\frac{-P_n^{\alpha,\beta}(-1)}{K_{n-1}(-1,-1)P_{n-1}^{\alpha,\beta+2}(-1)} = \frac{2^{\alpha+\beta+2}\Gamma(n)\Gamma(\beta+2)\Gamma(\beta+3)\Gamma(n+\alpha)}{\Gamma(n+\beta+2)\Gamma(n+\alpha+\beta+2)} = h_n(\alpha,\beta).$$

Also from (20)

$$\lim_{M\to\infty} M[x_{n,k+1}^M(\alpha,\beta) - x_{n-1,k}(\alpha,\beta+2)]$$
$$= \frac{-P_n^{\alpha,\beta}(x_{n-1,k}(\alpha,\beta+2))}{K_{n-1}(-1,-1)(x_{n-1,k}(\alpha,\beta+2)+1)[P_{n-1}^{\alpha,\beta+2}(x)]'\big|_{x=x_{n-1,k}(\alpha,\beta+2)}}.$$

On the other hand, it follows from

$$n(n+\alpha)(1+x)P_{n-1}^{\alpha,\beta+2}(x) = n(n+\alpha+\beta+1)P_n^{\alpha,\beta}(x) + (\beta+1)(1-x)[P_n^{\alpha,\beta}(x)]'$$

that

$$\begin{aligned} &n(n+\alpha+\beta+1)P_n^{\alpha,\beta}(x_{n-1,k}(\alpha,\beta+2)) \\ &\quad = -(\beta+1)(1-x_{n-1,k}(\alpha,\beta+2))[P_n^{\alpha,\beta}(x)]'\big|_{x=x_{n-1,k}(\alpha,\beta+2)} \end{aligned}$$

and

$$\begin{aligned} &n(n+\alpha)(1+x_{n-1,k}(\alpha,\beta+2))[P_n^{\alpha,\beta}(x)]'\big|_{x=x_{n-1,k}(\alpha,\beta+2)} \\ &\quad = [n(n+\alpha+\beta+1)-(\beta+1)][P_n^{\alpha,\beta}(x)]'\big|_{x=x_{n-1,k}(\alpha,\beta+2)} \\ &\qquad +(\beta+1)(1-x_{n-1,k}(\alpha,\beta+2))[P_n^{\alpha,\beta}(x)]''\big|_{x=x_{n-1,k}(\alpha,\beta+2)}. \end{aligned}$$

Now, using the last two equalities and the differential equation for the Jacobi polynomials

$$(1-x^2)[P_n^{\alpha,\beta}(x)]'' + [\beta-\alpha-(\alpha+\beta+1)x][P_n^{\alpha,\beta}(x)]' + n(n+\alpha+\beta+1)P_n^{\alpha,\beta}(x) = 0$$

we obtain

$$\begin{aligned} &(1+x_{n-1,k}(\alpha,\beta+2))[P_n^{\alpha,\beta+2}(x)]'\big|_{x=x_{n-1,k}(\alpha,\beta+2)} \\ &\quad = \frac{-(n+\beta+1)(n+\alpha+\beta+1)}{(\beta+1)(1-x_{n-1,k}(\alpha,\beta+2))}P_n^{\alpha,\beta}(x_{n-1,k}(\alpha,\beta+2)). \end{aligned}$$

Therefore

$$\begin{aligned} &\lim_{M\to\infty} M[x_{n,k+1}^M(\alpha,\beta) - x_{n-1,k}(\alpha,\beta+2)] \\ &= \frac{-P_n^{\alpha,\beta}(x_{n-1,k}(\alpha,\beta+2))}{K_{n-1}(-1,-1)(x_{n-1,k}(\alpha,\beta+2)+1)[P_{n-1}^{\alpha,\beta+2}(x)]'\big|_{x=x_{n-1,k}(\alpha,\beta+2)}} \\ &= \frac{\left[1-x_{n-1,k}(\alpha,\beta+2)\right]h_n(\alpha,\beta)}{2(\beta+2)}. \end{aligned}$$

Theorem 6 *The inequalities*

$$\begin{aligned} &x_{n,1}(\alpha,\beta) < x_{n,1}^N(\alpha,\beta) < x_{n-1,1}(\alpha+2,\beta) < \cdots < \\ &x_{n,n-1}(\alpha,\beta) < x_{n,n-1}^N(\alpha,\beta) < x_{n-1,n-1}(\alpha+2,\beta) < x_{n,n}(\alpha,\beta) < x_{n,n}^N(\alpha,\beta) < 1 \end{aligned}$$

hold for every $\alpha,\beta > -1$*. Moreover, each* $x_{n,k}^N(\alpha,\beta)$ *is an increasing function of* N *and, for each* $k = 1,\ldots,n-1$*,*

$$\lim_{N\to\infty} x_{n,n}^N(\alpha,\beta) = 1, \quad \lim_{N\to\infty} x_{n,k}^N(\alpha,\beta) = x_{n-1,k}(\alpha+2,\beta),$$

and

$$\lim_{N\to\infty} N[1-x_{n,n}^N(\alpha,\beta)] = g_n(\alpha,\beta),$$

$$\lim_{N\to\infty} N[x_{n-1,k}(\alpha+2,\beta) - x_{n,k}^N(\alpha,\beta)] = \frac{\left[1+x_{n-1,k}(\alpha+2,\beta)\right]g_n(\alpha,\beta)}{2(\alpha+2)},$$

where

$$g_n(\alpha,\beta)=\frac{2^{\alpha+\beta+2}\Gamma(n)\Gamma(\alpha+2)\Gamma(\alpha+3)\Gamma(n+\beta)}{\Gamma(n+\alpha+2)\Gamma(n+\alpha+\beta+2)}.$$

Proof The prove follows in the same way as the Theorem 5. We only observe that

$$\frac{n+\beta}{2n}(x-1)P_{n-1}^{\alpha+2,\beta}(x)=P_n^{\alpha,\beta}(x)-\frac{\alpha+1}{n(n+\alpha+\beta+1)}(1+x)[P_n^{\alpha,\beta}(x)]'.$$

In order to illustrate the results of Theorem 6, we enclose the graphs of $P_3^{\alpha,\beta,N+\varepsilon}(x)$, for $\alpha=\beta=0$ and some values of $\varepsilon>0$, in order to show the monotonicity of the zeros of $P_3^{\alpha,\beta,N}(x)$ as a function of the mass N (see Figure 1).



Fig. 1 The graphs of $P_3^{\alpha,\beta,N+\varepsilon}(x)$ for some values of ε.

In Table 1 we show the value of zeros of $P_3^{\alpha,\beta,N}(x)$, with $\alpha=\beta=0$, for several choices of N Notice that the largest zero converges to 1 and the other two zeros converge to the zeros of the Jacobi polynomial $P_2^{2,0}(x)$, that is, they converge to $x_{2,1}(2,0)=-0.75497$ and $x_{2,2}(4)=0.0883037$. Note also that all the zeros increase when N increase.

Table 1 Zeros of $P_3^{\alpha,\beta,N}(x)$ for some values of N.

N	$x_{3,1}^N(0,0)$	$x_{3,2}^N(0,0)$	$x_{3,3}^N(0,0)$
0	−0.774597	0	0.774597
1	−0.757872	0.0753429	0.955257
10	−0.755305	0.0868168	0.994575
100	−0.755004	0.0881528	0.999446
1000	−0.754974	0.0882886	0.999944

4.2 Laguerre type (Laguerre-Koornwinder) orthogonal polynomials

Next, we will deal with $p_n(x) = L_n^{\alpha}(x)$, that is, the classical monic Laguerre polynomials, which are orthogonal with respect to the measure $d\mu_{\alpha} = x^{\alpha}e^{-x}dx$, $\alpha > -1$, supported on $[0, +\infty)$. We will consider the Uvarov perturbation on μ_{α} with $a = 0$

$$d\mu_N = d\mu_{\alpha} + N\delta_0, N \geq 0. \tag{28}$$

The polynomials $L_n^{\alpha,N}(x)$, orthogonal with respect to (28), were also obtained by T. H. Koornwinder [14] as a special limit case of the Jacobi-Koornwinder (Jacobi type) orthogonal polynomials. In this direction, concerning analytic properties of these polynomials, many contributions have been done in the last years (see [1], [4], [7], [13], among others). The connection formula of $L_n^{\alpha,N}(x)$ is

$$L_n^{\alpha,N}(x) = L_n^{\alpha}(x) + NK_{n-1}(0,0)\,x\,L_{n-1}^{\alpha+2}(x), \tag{29}$$

where

$$K_{n-1}(0,0) = \frac{\Gamma(n+\alpha+1)}{\Gamma(n)\Gamma(\alpha+1)\Gamma(\alpha+2)}.$$

Now, we will analyze the behavior of their zeros. Let denote by $(x_{n,k}^N(\alpha))$ and $(x_{n,k}(\alpha))$ the zeros of the Laguerre type and the classical Laguerre orthogonal polynomials, respectively, arranged in an increasing order. Applying the results of Section 4, we obtain

Theorem 7 *The inequalities*

$$0 < x_{n,1}^N(\alpha) < x_{n,1}(\alpha) < x_{n-1,1}(\alpha+2) < x_{n,2}^N(\alpha) < x_{n,2}(\alpha) < \cdots$$
$$< x_{n-1,n-1}(\alpha+2) < x_{n,n}^N(\alpha) < x_{n,n}(\alpha)$$

hold for every $\alpha > -1$. Moreover, each $x_{n,k}^N(\alpha)$ is a decreasing function of N and, for each $k = 1, \ldots, n-1$,

$$\lim_{N\to\infty} x_{n,1}^N(\alpha) = 0, \quad \lim_{N\to\infty} x_{n,k+1}^N(\alpha) = x_{n-1,k}(\alpha+2),$$

as well as

$$\lim_{N\to\infty} Nx_{n,1}^N(\alpha) = g_n(\alpha),$$

$$\lim_{N\to\infty} N[x_{n,k+1}^N(\alpha) - x_{n-1,k}(\alpha+2)] = \frac{g_n(\alpha)}{\alpha+2},$$

where

$$g_n(\alpha) = \frac{\Gamma(n)\Gamma(\alpha+2)\Gamma(\alpha+3)}{\Gamma(n+\alpha+2)}. \tag{30}$$

Proof It remains to show the limits above. From (20)

$$\lim_{N\to\infty} Nx_{n,1}^N(\alpha) = \frac{-L_n^{\alpha}(0)}{K_{n-1}(0,0)L_{n-1}^{\alpha+2}(0)}.$$

Since

$$L_n^{\alpha}(0) = \frac{(-1)^n\Gamma(n+\alpha+1)}{\Gamma(\alpha+1)} \quad \text{and} \quad K_{n-1}(0,0) = \frac{\Gamma(n+\alpha+1)}{\Gamma(n)\Gamma(\alpha+1)\Gamma(\alpha+2)},$$

we obtain
$$\frac{-L_n^\alpha(0)}{K_{n-1}(0,0)L_{n-1}^{\alpha+2}(0)} = \frac{\Gamma(n)\Gamma(\alpha+2)\Gamma(\alpha+3)}{\Gamma(n+\alpha+2)} = g_n(\alpha).$$

Also from (20)

$$\lim_{N\to\infty} N[x_{n,k+1}^N(\alpha) - x_{n-1,k}(\alpha+2)]$$
$$= \frac{-L_n^\alpha(x_{n-1,k}(\alpha+2))}{K_{n-1}(0,0)x_{n-1,k}(\alpha+2)[L_{n-1}^{\alpha+2}(x)]'\big|_{x=x_{n-1,k}(\alpha+2)}}.$$

On the other hand, it is easily to verify that

$$xL_{n-1}^{\alpha+2}(x) = L_n^\alpha(x) + \frac{\alpha+1}{n}[L_n^\alpha(x)]'.$$

Thus,
$$[L_n^\alpha(x)]'\big|_{x=x_{n-1,k}(\alpha+2)} = -\frac{n}{\alpha+1}L_n^\alpha(x_{n-1,k}(\alpha+2))$$

and
$$x_{n-1,k}(\alpha+2)[L_n^\alpha(x)]'\big|_{x=x_{n-1,k}(\alpha+2)}$$
$$= [L_n^\alpha(x)]'\big|_{x=x_{n-1,k}(\alpha+2)} + \frac{\alpha+1}{n}[L_n^\alpha(x)]''\big|_{x=x_{n-1,k}(\alpha+2)}.$$

Now, using the last two equalities and the differential equation for the Laguerre polynomials
$$x[L_n^\alpha(x)]'' + (\alpha+1-x)[L_n^\alpha(x)]' + nL_n^\alpha(x) = 0$$

we obtain

$$x_{n-1,k}(\alpha+2)[L_n^\alpha(x)]'\big|_{x=x_{n-1,k}(\alpha+2)} = \frac{-(n+\alpha+1)}{\alpha+1}L_n^\alpha(x_{n-1,k}(\alpha+2)).$$

Therefore

$$\lim_{N\to\infty} N[x_{n,k+1}^N(\alpha) - x_{n-1,k}(\alpha+2)]$$
$$= \frac{-L_n^\alpha(x_{n-1,k}(\alpha+2))}{K_{n-1}(0,0)x_{n-1,k}(\alpha+2)[L_{n-1}^{\alpha+2}(x)]'\big|_{x=x_{n-1,k}(\alpha+2)}}$$
$$= \frac{\Gamma(n)\Gamma(\alpha+2)\Gamma(\alpha+2)}{\Gamma(n+\alpha+2)}$$
$$= \frac{g_n(\alpha)}{\alpha+2}.$$

In order to illustrate the results of Theorem 7, we enclose the graphs of $L_3^{\alpha,N+\varepsilon}(x)$, for $\alpha = 2$ and some values of $\varepsilon > 0$, in order to show the monotonicity of the zeros of $L_3^{\alpha,N}(x)$ as a function of the mass N. See Figure 2.

Table 2 shows the zeros of $L_3^{\alpha,N}(x)$, with $\alpha = 2$, for several choices of N. Observe that the smallest zero converges to 0 and the other two zeros converge to the zeros of the Laguerre polynomial $L_2^4(x)$, that is, they converge to $x_{2,1}(4) = 3.55051$ and $x_{2,2}(4) = 8.44949$. Notice that all the zeros decrease when N increases.



Fig. 2 The graphs of $L_3^{\alpha,N+\varepsilon}(x)$ for some values of ε.

Table 2 Zeros of $L_3^{\alpha,N}(x)$ for some values of N.

N	$x_{3,1}^N(2)$	$x_{3,2}^N(2)$	$x_{3,3}^N(2)$
0	1.51739	4.31158	9.17103
1	0.321731	3.64053	8.53774
10	0.0390611	3.5604	8.45936
100	0.00399042	3.55151	8.45049
1000	0.00039990	3.55061	8.44959

4.3 Hermite type orthogonal polynomials

Using the symmetrization process for the special case of Laguerre type orthogonal polynomials when $\alpha = -1/2$, we obtain the Hermite type orthogonal polynomials $H_n^N(x)$, which are orthogonal with respect to the measure

$$d\mu_N = e^{-x^2}dx + N\delta_0.$$

whose support is the real line. It easy to see that they are symmetric with respect to the origin and

$$H_{2n}^N(x) = L_n^{(-1/2,N)}(x^2) \tag{31}$$

and

$$H_{2n+1}^N(x) = H_{2n+1}^0(x) = xL_n^{(1/2,0)}(x^2). \tag{32}$$

Let denote by $(h_{2n,k}^N)$, $1 \leq k \leq 2n$, and $(h_{2n+1,k})$, $1 \leq k \leq 2n+1$, the zeros of the Hermite type polynomials $H_{2n}^N(x)$ and $H_{2n+1}^N(x)$, respectively, ordered as follows : $h_{2n,n}^N < \cdots < h_{2n,1}^N(\alpha)$ and $h_{2n+1,n} < \cdots < h_{2n+1,1}$. Because of the symmetry property of these polynomials, we get $h_{2n,k}^N = -h_{2n,2n-k+1}^N$ and $h_{2n+1,k} = -h_{2n+1,2n+2-k}$, $k = 1, \ldots, n$. Furthermore, from (31) and (32), we have

$$\left[h_{2n,k}^N\right]^2 = x_{n,n-k+1}^N(-1/2)$$

and

$$\left[h_{2n+1,k}\right]^2 = x_{n,n-k+1}(1/2).$$

Then, as a straightforward consequence of Theorem 7, we get

Theorem 8 *Let $n \in \mathbb{N}$. Then*

(i) The inequalities

$$0 < \left[h^N_{2n,n}\right]^2 < x_{n,1}(-1/2) < x_{n-1,1}(3/2) < \left[h^N_{2n,n-1}\right]^2 < x_{n,2}(-1/2) < \cdots$$
$$< x_{n-1,n-1}(3/2) < \left[h^N_{2n,1}\right]^2 < x_{n,n}(-1/2)$$

hold. Moreover, each $h^N_{2n,k}$, $k = 1, \ldots, n$, is a decreasing function of N and, for each $k = 1, \ldots, n-1$,

$$\lim_{N\to\infty} \left[h^N_{2n,n}\right]^2 = 0, \quad \lim_{N\to\infty} \left[h^N_{2n,k}\right]^2 = x_{n-1,n-k}(3/2),$$

and

$$\begin{aligned} &\lim_{N\to\infty} N \left[h^N_{2n,n}\right]^2 = g_n(-1/2), \\ &\lim_{N\to\infty} N[\left[h^N_{2n,k}\right]^2 - x_{n-1,n-k}(3/2)] = \frac{g_n(-1/2)}{3/2}; \end{aligned} \tag{33}$$

(ii) The inequalities

$$0 < \left[h_{2n+1,n}\right]^2 < x_{n,1}(1/2) < x_{n-1,1}(5/2) < \left[h_{2n+1,n-1}\right]^2 < x_{n,2}(1/2) < \cdots$$
$$< x_{n-1,n-1}(5/2) < \left[h_{2n,1}\right]^2 < x_{n,n}(1/2)$$

hold.

The function $g_n(\alpha)$ in (33) is defined in (30). More details about the zeros of Hermite-type polynomials can be found in [18].

5 Electrostatic interpretation

5.1 Some preliminaries results

We also provided an alternative connection formula for the MOPS $\{p^N_n(a;x)\}_{n\geq 0}$ in terms of the kernel polynomials $\{p^*_n(a;x)\}_{n\geq 0}$.

Theorem 9 *The polynomials $\{p^N_n(a;x)\}_{n\geq 0}$ can be also represented as*

$$p^N_n(a;x) = p^*_n(a;x) + c_n\, p^*_{n-1}(a;x), \tag{34}$$

where

$$c_n = -\frac{1 + NK_n(a,a)}{1 + NK_{n-1}(a,a)} \frac{p_{n-1}(a)}{p_n(a)} \gamma_n \quad \text{and} \quad \gamma_n = \frac{\|p_n\|^2_\mu}{\|p_{n-1}\|^2_\mu}. \tag{35}$$

Proof Using (4), we can write $p_n(x)$ as

$$p_n(x) = \frac{\|p_n\|_\mu^2}{p_n(a)} \left[K_n(a,x) - K_{n-1}(a,x)\right]. \tag{36}$$

Now, from (7) and (36), we have

$$p_n(x) = p_n^*(a;x) - \gamma_n \frac{p_{n-1}(a)}{p_n(a)} p_{n-1}^*(a;x). \tag{37}$$

Therefore, substituting (7) and (37) in (15), we obtain

$$\begin{aligned} p_n^N(a;x) &= \left(p_n^*(a;x) - \gamma_n \frac{p_{n-1}(a)}{p_n(a)} p_{n-1}^*(a;x) \right) \\ &\qquad - \frac{N p_n(a)}{1 + N K_{n-1}(a,a)} \frac{p_{n-1}(a)}{\|p_{n-1}\|_\mu^2} p_{n-1}^*(a;x) \\ &= p_n^*(a;x) + c_n\, p_{n-1}^*(a;x), \end{aligned}$$

where c_n is given in (35).

Theorem 10 *The MOPS $\{p_n^*(a;x)\}_{n\geq 0}$ satisfies the three-term recurrence relation*

$$x\, p_n^*(a;x) = p_{n+1}^*(a;x) + \beta_n^*\, p_n^*(a;x) + \gamma_n^*\, p_{n-1}^*(a;x), \quad n \geq 0, \tag{38}$$

with initial conditions $p_0^*(a;x) = 1$ and $p_{-1}^*(a;x) = 0$. The coefficients of the TTRR are

$$\beta_n^* = \beta_{n+1} + \frac{p_{n+2}(a)}{p_{n+1}(a)} - \frac{p_{n+1}(a)}{p_n(a)}, \quad n \geq 0, \tag{39}$$

and

$$\gamma_n^* = \frac{p_{n+1}(a) p_{n-1}(a)}{[p_n(a)]^2} \gamma_n > 0 \quad n \geq 1. \tag{40}$$

Proof From (7) and (37), we have

$$\begin{aligned} (x-a) p_n^*(a;x) &= p_{n+1}(x) - \frac{p_{n+1}(a)}{p_n(a)} p_n(x) \\ &= p_{n+1}^*(a;x) - \gamma_{n+1} \frac{p_n(a)}{p_{n+1}(a)} p_n^*(a;x) \\ &\qquad - \frac{p_{n+1}(a)}{p_n(a)} \left(p_n^*(a;x) - \gamma_n \frac{p_{n-1}(a)}{p_n(a)} p_{n-1}^*(a;x) \right) \\ &= p_{n+1}^*(a;x) - \left(\gamma_{n+1} \frac{p_n(a)}{p_{n+1}(a)} + \frac{p_{n+1}(a)}{p_n(a)} \right) p_n^*(a;x) \\ &\qquad + \gamma_n \frac{p_{n+1}(a) p_{n-1}(a)}{[p_n(a)]^2} p_{n-1}^*(a;x), \end{aligned}$$

or, equivalently,

$$\begin{aligned} x p_n^*(a;x) &= p_{n+1}^*(a;x) - \left(\gamma_{n+1} \frac{p_n(a)}{p_{n+1}(a)} + \frac{p_{n+1}(a)}{p_n(a)} - a \right) p_n^*(a;x) \\ &\qquad + \gamma_n \frac{p_{n+1}(a) p_{n-1}(a)}{[p_n(a)]^2} p_{n-1}^*(a;x). \end{aligned}$$

Since

$$ap_{n+1}(a) = p_{n+2}(a) + \beta_{n+1}p_{n+1}(a) + \gamma_{n+1}p_n(a),$$

we obtain

$$\gamma_{n+1}p_n(a) = ap_{n+1}(a) - p_{n+2}(a) - \beta_{n+1}p_{n+1}(a).$$

Thus,

$$\begin{aligned}
\beta_n^* &= -\left(\gamma_{n+1}\frac{p_n(a)}{p_{n+1}(a)} + \frac{p_{n+1}(a)}{p_n(a)}\right) + a \\
&= -\left(\frac{ap_{n+1}(a) - p_{n+2}(a) - \beta_{n+1}p_{n+1}(a)}{p_{n+1}(a)} + \frac{p_{n+1}(a)}{p_n(a)}\right) + a \\
&= -\frac{ap_{n+1}(a)}{p_{n+1}(a)} + \frac{p_{n+2}(a)}{p_{n+1}(a)} + \frac{\beta_{n+1}p_{n+1}(a)}{p_{n+1}(a)} - \frac{p_{n+1}(a)}{p_n(a)} + a \\
&= \beta_{n+1} + \frac{p_{n+2}(a)}{p_{n+1}(a)} - \frac{p_{n+1}(a)}{p_n(a)}.
\end{aligned}$$

For the other coefficient, we have

$$\gamma_n^* = \frac{\|p_n^*\|_{\mu^*}^2}{\|p_{n-1}^*\|_{\mu^*}^2} = \frac{\langle (x-a)\,p_n^*, p_n^*\rangle_\mu}{\left\langle (x-a)\,p_{n-1}^*, p_{n-1}^*\right\rangle_\mu}. \tag{41}$$

But according to (7)

$$(x-a)\,p_n^*(a;x) = p_{n+1}(x) - \frac{p_{n+1}(a)}{p_n(a)}p_n(x),$$

and

$$\left\langle (x-a)\,p_n^*\,(a;x)\,, p_n^*\,(a;x)\right\rangle_\mu = -\frac{p_{n+1}(a)}{p_n(a)}\left\|p_n^*\right\|_\mu^2.$$

Thus (41) becomes

$$\gamma_n^* = \frac{p_{n+1}(a)p_{n-1}(a)}{[p_n(a)]^2}\gamma_n.$$

As an example, we can analyze the behavior of these coefficients in the Laguerre case. Let $\widehat{L}_n^\alpha\,(x) = \frac{(-1)^n}{n!}L_n^\alpha\,(x)$, then we have

$$\begin{aligned}
\frac{\beta_n^*}{\beta_{n+1}} &= 1 - \frac{1}{\beta_{n+1}}\,(n+2)\,\frac{\widehat{L}_{n+2}^\alpha\,(a)}{\widehat{L}_{n+1}^\alpha\,(a)} + \frac{1}{\beta_{n+1}}\,(n+1)\,\frac{\widehat{L}_{n+1}^\alpha\,(a)}{\widehat{L}_n^\alpha\,(a)} \\
&= 1 - \frac{n+2}{2n+\alpha+3}\left(1 + \frac{\sqrt{|a|}}{\sqrt{n+2}} + \mathcal{O}\left(\frac{1}{n+2}\right)\right) \\
&\quad + \frac{n+1}{2n+\alpha+3}\left(1 + \frac{\sqrt{|a|}}{\sqrt{n+1}} + \mathcal{O}\left(\frac{1}{n+1}\right)\right) \\
&= 1 - \frac{1}{2n+\alpha+3} + \frac{\sqrt{|a|}}{2n+\alpha+3}\left(-\sqrt{n+2} + \sqrt{n+1}\right) + \mathcal{O}\left(\frac{1}{n^2}\right) \\
&= 1 - \frac{1}{2n+\alpha+3} - \frac{\sqrt{|a|}}{2n+\alpha+3}\frac{1}{\sqrt{n+2}+\sqrt{n+1}} + \mathcal{O}\left(\frac{1}{n^2}\right) \\
&= 1 - \frac{1}{2n+\alpha+3} - \frac{\sqrt{|a|}}{2}n^{-3/2} + \mathcal{O}\left(\frac{1}{n^2}\right).
\end{aligned}$$

Thus

$$\begin{aligned}
\frac{\beta_n^*}{\beta_n} &= \frac{2n+\alpha+3}{2n+\alpha+1}\left(1-\frac{1}{2n+\alpha+3}-\frac{\sqrt{|a|}}{2}n^{-3/2}+\mathcal{O}\left(\frac{1}{n^2}\right)\right)\\
&= \left(1+\frac{2}{2n+\alpha+1}\right)\left(1-\frac{1}{2n+\alpha+3}-\frac{\sqrt{|a|}}{2}n^{-3/2}+\mathcal{O}\left(\frac{1}{n^2}\right)\right)\\
&= 1+\frac{1}{2n}+\mathcal{O}\left(n^{-3/2}\right).
\end{aligned}$$

On the other hand, taking into account

$$\gamma_n^* = \frac{p_{n+1}(a)p_{n-1}(a)}{\left[p_n(a)\right]^2}\gamma_n,$$

then

$$\begin{aligned}
\frac{\gamma_n^*}{\gamma_n} &= \frac{(n+1)}{n}\frac{\widehat{L}_{n+1}^{\alpha}(a)}{\widehat{L}_{n}^{\alpha}(a)}\frac{\widehat{L}_{n-1}^{\alpha}(a)}{\widehat{L}_{n}^{\alpha}(a)}\\
&= \left(1+\frac{1}{n}\right)\frac{1+\frac{\sqrt{|a|}}{\sqrt{n+1}}+\mathcal{O}\left(\frac{1}{n+1}\right)}{1+\frac{\sqrt{|a|}}{\sqrt{n}}+\mathcal{O}\left(\frac{1}{n}\right)}\\
&= \left(1+\frac{1}{n}\right)\left(1+\frac{\sqrt{|a|}}{\sqrt{n+1}}+\mathcal{O}\left(\frac{1}{n+1}\right)\right)\left(1-\frac{\sqrt{|a|}}{\sqrt{n}}+\mathcal{O}\left(\frac{1}{n}\right)\right)\\
&= \left(1+\frac{1}{n}\right)\left(1-\frac{\sqrt{|a|}}{2n^{3/2}}+\mathcal{O}\left(\frac{1}{n^2}\right)\right)\\
&= 1+\frac{1}{n}+\mathcal{O}\left(n^{-3/2}\right).
\end{aligned}$$

Next, we will assume that $d\mu^*(x) = (x-a)\,\omega(x)\,dx$ where $\omega(x)$ is a weight function supported on the real line. We can associate with $\omega(x)$ an external potential $\upsilon(x)$ such that $\omega(x) = \exp(-\upsilon(x))$.

Notice that if $\upsilon(x)$ is assumed to be differentiable in the support of $d\mu(x) = \omega(x)\,dx$ then

$$\frac{\omega'(x)}{\omega(x)} = -\upsilon'(x).$$

If $\upsilon'(x)$ is a rational function, then the weight function $\omega(x)$ is said to be semi-classical (see [17], [20]). The linear functional u associated with $\omega(x)$, i.e.,

$$\langle u, p(x)\rangle = \int_{\Sigma} p(x)\omega(x)\,dx,$$

satisfies a distributional equation (which is known in the literature as Pearson equation)

$$D(\sigma(x)u) = \tau(x)u,$$

where $\sigma(x)$ and $\tau(x)$ are polynomials such that $\sigma(x)$ is monic and $\deg(\tau(x)) \geq 1$.

Notice that, in terms of the weight function, the above relation means that

$$\frac{\omega'(x)}{\omega(x)} = \frac{\tau(x) - \sigma'(x)}{\sigma(x)},$$

or, equivalently,

$$\upsilon'(x) = -\frac{\tau(x) - \sigma'(x)}{\sigma(x)}.$$

Let consider the linear functional u^* associated with the measure $\mu^*(x)$. In order to find the Pearson equation that u^* satisfies we will analyze two situations.

(i) If $\sigma(a) \neq 0$, then

$$\begin{aligned} D\left((x-a)\sigma(x)u^*\right) &= D\left((x-a)^2\sigma(x)u\right) \\ &= 2(x-a)\sigma(x)u + (x-a)^2 D(\sigma(x)u) \\ &= 2\sigma(x)u^* + (x-a)^2\tau(x)u \\ &= \left[2\sigma(x) + (x-a)\tau(x)\right]u^*. \end{aligned}$$

Thus,

$$D\left(\phi(x)u^*\right) = \psi(x)u^*,$$

where

$$\left| \begin{aligned} &\phi(x) = (x-a)\sigma(x) \\ &\psi(x) = 2\sigma(x) + (x-a)\tau(x). \end{aligned} \right. \tag{42}$$

(ii) If $\sigma(a) = 0$, i.e., $\sigma(x) = (x-a)\tilde{\sigma}(x)$, then

$$\begin{aligned} D\left(\sigma(x)u^*\right) &= D\left((x-a)\tilde{\sigma}(x)u^*\right) \\ &= D\left((x-a)^2\tilde{\sigma}(x)u\right) = D\left((x-a)\sigma(x)u\right) \\ &= \sigma(x)u + (x-a)D\left(\sigma(x)u\right) = \sigma(x)u + (x-a)\tau(x)u \\ &= \left(\tilde{\sigma}(x) + \tau(x)\right)u^*. \end{aligned}$$

In this case,

$$D\left(\phi(x)u^*\right) = \psi(x)u^*,$$

with

$$\left| \begin{aligned} &\phi(x) = \sigma(x) \\ &\psi(x) = \tilde{\sigma}(x) + \tau(x). \end{aligned} \right. \tag{43}$$

It is a very well known result that the sequence of monic polynomials $\{p_n^*(a;x)\}_{n\geq 0}$, orthogonal with respect to u^* satisfies a structure relation (see [6] and [17])

$$\phi(x)D\left(p_n^*(a;x)\right) = A(x,n)p_n^*(a;x) + B(x,n)p_{n-1}^*(a;x) \tag{44}$$

where $A(x,n)$ and $B(x,n)$ are polynomials of fixed degree, that do not depend on n.

Lemma 2 *[9] We have*

$$A(x,n)+A(x,n-1)+\frac{(x-\beta^*_{n-1})}{\gamma^*_{n-1}}B(x,n-1)=\phi'(x)-\psi(x). \tag{45}$$

Proof According to a result by Ismail ([9], (1.12)) which must be adapted to our situation since we use monic polynomials, we get

$$\begin{aligned}A(x,n)+A(x,n-1)+\frac{(x-\beta^*_{n-1})}{\gamma^*_{n-1}}B(x,n-1)&=-\phi(x)\frac{[\omega^*(x)]'}{\omega^*(x)}\\&=-\phi(x)\frac{\psi(x)-\phi'(x)}{\phi(x)}\\&=\phi'(x)-\psi(x),\end{aligned}$$

where $\omega^*(x)=(x-a)\omega(x)$.

Now, applying the derivative operator in (34) and multiplying it by $\phi(x)$, we obtain

$$\phi(x)D\left(p_n^N(a;x)\right)=\phi(x)D\left(p_n^*(a;x)\right)+c_n\phi(x)D\left(p^*_{n-1}(a;x)\right). \tag{46}$$

Thus, substituting (44) in (46), yields

$$\begin{aligned}\phi(x)D\left(p_n^N(a;x)\right)=A(x,n)p_n^*(a;x)+\left[B(x,n)+c_nA(x,n-1)\right]p^*_{n-1}(a;x)\\+c_nB(x,n-1)p^*_{n-2}(a;x).\end{aligned} \tag{47}$$

Finally, using the TTRR (38) in (47), we obtain

$$\phi(x)\left(p_n^N(a;x)\right)'=A^*(x,n)p_n^*(a;x)+B^*(x,n)p^*_{n-1}(a;x), \tag{48}$$

where

$$A^*(x,n)=\left(A(x,n)-\frac{c_n}{\gamma^*_{n-1}}B(x,n-1)\right) \tag{49}$$

and

$$B^*(x,n)=\left(B(x,n)+c_nA(x,n-1)+\frac{c_n}{\gamma^*_{n-1}}(x-\beta^*_{n-1})B(x,n-1)\right). \tag{50}$$

Therefore, from (34) and (48), it follows that

$$\begin{bmatrix}1 & c_n\\ A^*(x,n) & B^*(x,n)\end{bmatrix}\begin{bmatrix}p_n^*(a;x)\\ p^*_{n-1}(a;x)\end{bmatrix}=\begin{bmatrix}p_n^N(a;x)\\ \phi(x)D\left(p_n^N(a;x)\right)\end{bmatrix},$$

that is,

$$p_n^*(a;x)=\frac{\begin{vmatrix}p_n^N(a;x) & c_n\\ \phi(x)D\left(p_n^N(a;x)\right) & B^*(x,n)\end{vmatrix}}{\begin{vmatrix}1 & c_n\\ A^*(x,n) & B^*(x,n)\end{vmatrix}}$$

and

$$p_{n-1}^{*}(a;x)=\frac{\begin{vmatrix}1 & p_n^N(a;x)\\ A^{*}(x,n) & \phi(x)D\left(p_n^N(a;x)\right)\end{vmatrix}}{\begin{vmatrix}1 & c_n\\ A^{*}(x,n) & B^{*}(x,n)\end{vmatrix}},$$

or, equivalently,

$$p_n^{*}(a;x)=\frac{B^{*}(x,n)}{B^{*}(x,n)-c_nA^{*}(x,n)}\,p_n^N(a;x)-\frac{c_n\phi(x)}{B^{*}(x,n)-c_nA^{*}(x,n)}\,D\left(p_n^N(a;x)\right) \tag{51}$$

and

$$p_{n-1}^{*}(a;x)=\frac{-A^{*}(x,n)}{B^{*}(x,n)-c_nA^{*}(x,n)}p_n^N(a;x)+\frac{\phi(x)}{B^{*}(x,n)-c_nA^{*}(x,n)}D\left(p_n^N(a;x)\right). \tag{52}$$

Now, substituting (51) and (52) in (44), we deduce

$$\begin{aligned}&\phi(x)D\left(\frac{B^{*}(x,n)}{B^{*}(x,n)-c_nA^{*}(x,n)}\,p_n^N(a;x)-\frac{c_n\phi(x)}{B^{*}(x,n)-c_nA^{*}(x,n)}\,D\left(p_n^N(a;x)\right)\right)\\ &=A(x,n)\left(\frac{B^{*}(x,n)}{B^{*}(x,n)-c_nA^{*}(x,n)}\,p_n^N(a;x)-\frac{c_n\phi(x)}{B^{*}(x,n)-c_nA^{*}(x,n)}\,D\left(p_n^N(a;x)\right)\right)\\ &+B(x,n)\left(\frac{-A^{*}(x,n)}{B^{*}(x,n)-c_nA^{*}(x,n)}p_n^N(a;x)+\frac{\phi(x)}{B^{*}(x,n)-c_nA^{*}(x,n)}D\left(p_n^N(a;x)\right)\right).\end{aligned}$$

Then a straightforward calculation yields

Theorem 11 *The MOPS* $\{p_n^N(a;x)\}_{n\geq 0}$ *satisfies the second order linear differential equation*

$$\mathcal{A}(x;n)(p_n^N(a;x))''+\mathcal{B}(x;n)(p_n^N(a;x))'+\mathcal{C}(x;n)p_n^N(a;x)=0, \tag{53}$$

where

$$\begin{aligned}\mathcal{A}(x;n)&=\frac{c_n\left[\phi(x)\right]^2}{B^{*}(x,n)-c_nA^{*}(x,n)},\\ \mathcal{B}(x;n)&=\frac{\phi(x)\left[B(x,n)-B^{*}(x,n)+c_n(\phi'(x)-A(x,n))\right]}{B^{*}(x,n)-c_nA^{*}(x,n)}\\ &\qquad-\frac{c_n\phi(x)^2\left(B^{*}(x,n)-c_nA^{*}(x,n)\right)'}{\left(B^{*}(x,n)-c_nA^{*}(x,n)\right)^2}\\ \mathcal{C}(x;n)&=\frac{A(x,n)B^{*}(x,n)-B(x,n)A^{*}(x,n)}{B^{*}(x,n)-c_nA^{*}(x,n)}-\phi(x)D\left(\frac{B^{*}(x,n)}{B^{*}(x,n)-c_nA^{*}(x,n)}\right).\end{aligned}$$

A different approach to this differential equation appears in [15] using the fact that the Uvarov transform of a semiclassical linear functional is again a semiclassical linear functional.

It is important to notice that for the electrostatic interpretation of the zeros is enough to consider the polynomial coefficients of $(p_n^N(a;x))''$ and $(p_n^N(a;x))'$. In fact, it will come from the ratio

$$\frac{(p_n^N(a;x))''}{(p_n^N(a;x))'} = -\frac{\mathcal{B}(x;n)}{\mathcal{A}(x;n)}$$

evaluated at the zeros of $p_n^N(a;x)$.

Let $(x_{n,k}^N)$ be the zeros of $p_n^N(a;x)$. If we evaluate the second-order linear differential equation (53) at $x_{n,k}^N$ then we obtain

$$\mathcal{A}(x_{n,k}^N;n)(p_n^N(a;x_{n,k}^N))'' + \mathcal{B}(x_{n,k}^N;n)(p_n^N(a;x_{n,k}^N))' = 0.$$

when $x_{n,k}^N$ is a zero of the polynomial $p_n^N(a;x)$.

Hence,

$$\frac{(p_n^N(a;x_{n,k}^N))''}{(p_n^N(a;x_{n,k}^N))'} = -\frac{\mathcal{B}(x_{n,k}^N;n)}{\mathcal{A}(x_{n,k}^N;n)}. \tag{54}$$

Substituting $\mathcal{A}(x_{n,k}^N;n)$ and $\mathcal{B}(x_{n,k}^N;n)$ in the right hand side of (54), we get

$$\begin{aligned}&\frac{(p_n^N(a;x_{n,k}^N))''}{(p_n^N(a;x_{n,k}^N))'}\\ &= \frac{(B^*(x_{n,k}^N,n) - c_nA^*(x_{n,k}^N,n))'}{B^*(x_{n,k}^N,n) - c_nA^*(x_{n,k}^N,n)} + \frac{B^*(x_{n,k}^N,n) - B(x_{n,k}^N,n) + c_nA(x_{n,k}^N,n) - c_n\phi'(x_{n,k}^N)}{c_n\phi(x_{n,k}^N)}.\end{aligned}$$

If we denote $Q(x) := B^*(x,n) - c_nA^*(x,n)$ and using (45), (49), and (50), then we have

$$\begin{aligned}Q(x) &= B^*(x,n) - c_nA^*(x,n)\\ &= B(x,n) + c_n\left[A(x,n-1) + \frac{x-\beta_{n-1}^*}{\gamma_{n-1}^*}B(x,n-1) - A(x,n) + \frac{c_n}{\gamma_{n-1}^*}B(x,n-1)\right]\\ &= B(x,n) + c_n\Bigg[-2A(x,n) + \left(A(x,n) + A(x,n-1) + \frac{(x-\beta_{n-1}^*)}{\gamma_{n-1}^*}B(x,n-1)\right)\\ &\qquad\qquad + \frac{c_n}{\gamma_{n-1}^*}B(x,n-1)\Bigg]\\ &= B(x,n) + c_n\left[-2A(x,n) + \phi'(x) - \psi(x) + \frac{c_n}{\gamma_{n-1}^*}B(x,n-1)\right],\end{aligned}$$

i.e.,

$$Q(x) = B(x,n) + c_n\left[-2A(x,n) + \phi'(x) - \psi(x) + \frac{c_n}{\gamma_{n-1}^*}B(x,n-1)\right]. \tag{55}$$

On the other hand, from (55) and (49), we obtain

$$\begin{aligned}
&B^*(x,n) - B(x,n) + c_n A(x,n) - c_n\phi'(x)\\
&= Q(x) + c_n A^*(x,n) - B(x,n) + c_n A(x,n) - c_n\phi'(x)\\
&= B(x,n) + c_n\left[-2A(x,n) + \phi'(x) - \psi(x) + \frac{c_n}{\gamma^*_{n-1}}B(x,n-1)\right]\\
&+c_n\left(A(x,n) - \frac{c_n}{\gamma^*_{n-1}}B(x,n-1)\right) - B(x,n) + c_n A(x,n) - c_n\phi'(x)\\
&= -c_n\psi(x).
\end{aligned}$$

Thus

$$\frac{(p_n^N(a;x^N_{n,k}))''}{(p_n^N(a;x^N_{n,k}))'} = D\left[\ln Q(x)\right]|_{x=x^N_{n,k}} - \frac{\psi(x^N_{n,k})}{\phi(x^N_{n,k})}. \tag{56}$$

We consider two external fields

$$-\int\frac{\psi(x)}{\phi(x)}dx \quad \text{and} \quad \ln Q(x),$$

in such a way that the total external potential $V(x)$ is given by

$$V(x) = -\int\frac{\psi(x)}{\phi(x)}dx + \ln Q(x). \tag{57}$$

Let introduce a system of n movable unit charges in $[a,\eta]$ or $[\xi,a]$, depending on the location of the point a with respect to $C_0(\Sigma) = [\xi,\eta]$, in the presence of the external potential $V(x)$ of (57). Let

$$x := (x_1,\dots,x_n),$$

where $x_1,\dots,x_n$ denote the positions of the particles. The total energy of the system is

$$E(x) = \sum_{k=1}^{n} V(x_k) - 2\sum_{1\le j<k\le n}\ln|x_j - x_k|. \tag{58}$$

Let

$$T(x) := \exp(-E(x)) = \left[\prod_{j=1}^{n}\frac{\exp(-\int\frac{\psi(x_j)}{\phi(x_j)}dx)}{Q(x_j)}\right]\prod_{1\le j<k\le n}(x_j - x_k)^2. \tag{59}$$

In order to find the critical points of $E(x)$ we will analyze the gradient of $\ln T(x)$. Indeed,

$$\frac{\partial}{\partial x_j}\ln T(x) = 0, \quad j = 1,\dots,n,$$

i.e,

$$-\frac{\partial}{\partial x_j}E(x) = 0 \Leftrightarrow \frac{\psi(x_j)}{\phi(x_j)} - \frac{Q'(x_j)}{Q(x_j)} + 2\sum_{1\le k\le n, k\ne j}\frac{1}{x_j - x_k} = 0, \quad j = 1,\dots,n. \tag{60}$$

Let

$$f(y) := (y - x_1)\cdots(y - x_n).$$

Thus,

$$\frac{\psi(x_j)}{\phi(x_j)} - \frac{Q'(x_j)}{Q(x_j)} + \frac{f''(x_j)}{f'(x_j)} = 0, \quad j = 1, \ldots, n,$$

or, equivalently,

$$f''(y) + \frac{\mathcal{B}(y;n)}{\mathcal{A}(y;n)} f'(y) = 0, \quad y = x_1, \ldots, x_n.$$

Therefore

$$f''(y) + \frac{\mathcal{B}(y;n)}{\mathcal{A}(y;n)} f'(y) + \frac{\mathcal{C}(y;n)}{\mathcal{A}(y;n)} f(y) = 0, \quad y = x_1, \ldots, x_n. \tag{61}$$

On the other hand, from (53) and (61) we get

$$f(y) = p_n^N(a;y),$$

and then the zeros of $p_n^N(a;y)$ satisfy (60).

5.2 Electrostatic interpretation of the zeros of Laguerre type orthogonal polynomials

Firstly, we shall enumerate some useful basic properties of the Laguerre classical monic polynomials $L_n^\alpha(x)$.

i. Let u be the linear funcional

$$\langle u, p \rangle = \int_0^{+\infty} p(x) x^\alpha e^{-x} dx, \quad \alpha > -1, \ p \in \mathbb{P}.$$

So u satisfies the Pearson differential equation

$$D(\sigma(x)u) = \tau(x)u,$$

where

$$\sigma(x) = x, \quad \tau(x) = \alpha + 1 - x. \tag{62}$$

ii. For every $n \in \mathbb{N}$,

$$\begin{aligned} &L_{-1}^\alpha(x) = 0, \quad L_0^\alpha(x) = 1, \\ &L_{n+1}^\alpha(x) = (x - \beta_n) L_n^\alpha(x) - \gamma_n L_{n-1}^\alpha(x), \end{aligned} \tag{63}$$

where

$$\beta_n = \beta_n^\alpha = 2n + \alpha + 1, \quad \gamma_n = \gamma_n^\alpha = n\,(n + \alpha)\,.$$

iii. For every $n \in \mathbb{N}$

$$L_n^\alpha(0) = (-1)^n \frac{\Gamma(n + \alpha + 1)}{\Gamma(\alpha + 1)}. \tag{64}$$

iv. For every $n \in \mathbb{N}$

$$K_n(0,0) = \frac{1}{n!} \frac{\Gamma(n + \alpha + 2)}{\Gamma(\alpha + 2)\Gamma(\alpha + 1)}. \tag{65}$$

v. For every $n \in \mathbb{N}$

$$x \left[L_n^\alpha(x)\right]' = n L_n^\alpha(x) + n(n + \alpha) L_{n-1}^\alpha(x). \tag{66}$$

Now, we shall give an electrostatic interpretation for the zeros of Laguerre type polynomials $L_n^{\alpha,N}(a;x)$ which are orthogonal with respect to the measure $d\mu_N = x^{\alpha}e^{-x}dx + N\delta_a$, that is, they are orthogonal with respect to the following inner product:

$$\langle p,q\rangle = \int_0^{+\infty} p(x)q(x)x^{\alpha}e^{-x}dx + Np(a)q(a), \quad a \le 0.$$

We will analyze two cases:

1. Firstly, we consider $a = 0$. Thus, the polynomials $L_n^{\alpha,N}(0;x)$ are orthogonal with respect to

$$d\mu_N = x^{\alpha}e^{-x}dx + N\delta_0.$$

Now, observe that the polynomials $p_n^*(0;x) = L_n^{\alpha+1}(x)$ associated with the measure

$$d\mu^*(x) = x^{\alpha+1}e^{-x}dx$$

have Pearson's coefficients given by (see (43) and (62))

$$\phi(x) = \sigma(x) = x, \quad \psi(x) = \widetilde{\sigma}(x) + \tau(x) = \alpha + 2 - x.$$

On the other hand, from (66), the structure relation (44) reads

$$\phi(x)D\left(L_n^{\alpha+1}(x)\right) = A(x,n)L_n^{\alpha+1}(x) + B(x,n)L_{n-1}^{\alpha+1}(x),$$

where

$$\phi(x) = x, \quad A(x,n) = n, \quad B(x,n) = n + \alpha + 1.$$

In this case, the coefficients (35) and (38) are given by

$$\begin{aligned} \gamma_n^* &= n(n+\alpha+1), \\ c_n &= -\frac{1 + NK_n(0,0)}{1 + NK_{n-1}(0,0)} \frac{L_{n-1}^{\alpha}(0)}{L_n^{\alpha}(0)} n(n+\alpha). \end{aligned}$$

Using (64) and (65), we obtain

$$\begin{aligned} c_n &= \frac{1 + N\dfrac{\Gamma(n+\alpha+2)}{n!\Gamma(\alpha+1)\Gamma(\alpha+2)}}{1 + N\dfrac{\Gamma(n+\alpha+1)}{(n-1)!\Gamma(\alpha+1)\Gamma(\alpha+2)}} \cdot n \\ &= \frac{n!\Gamma(\alpha+1)\Gamma(\alpha+2) + N\Gamma(n+\alpha+2)}{(n-1)!\Gamma(\alpha+1)\Gamma(\alpha+2) + N\Gamma(n+\alpha+1)}. \end{aligned}$$

As a conclusion, $Q(x)$ in (55) becomes

$$\begin{aligned} Q\left(x\right) &= B(x,n) + c_n\left[-2A(x,n) + \phi'(x) - \psi(x) + \frac{c_n}{\gamma_{n-1}^*}B(x,n-1)\right] \\ &= n(n+\alpha+1) + c_n\left[-2n + 1 - (\alpha + 2 - x) + c_n\right] \\ &= n(n+\alpha+1) - c_n\left(2n + 1 + \alpha - c_n\right) + c_n x \end{aligned}$$

and its zero will be denoted by

$$u_n = (2n + 1 + \alpha - c_n) - \frac{n(n + \alpha + 1)}{c_n}. \tag{67}$$

Now, it is easily to see that $0 < c_n < n + \alpha + 1$. Thus, $Q(0) < 0$ and it implies that $u_n > 0$.

Taking into account

$$\Gamma(z) \sim \sqrt{\frac{2\pi}{z}} e^{-z} z^z = \sqrt{2\pi} e^{-z} z^{z-\frac{1}{2}},$$

it is easy to see that

$$\begin{aligned} \frac{\Gamma(n+\alpha+1)}{\Gamma(n+1)} &\sim \frac{\sqrt{2\pi} e^{-n-1} (n+\alpha+1)^{n+\frac{1}{2}} (n+\alpha+1)^{\alpha} e^{-\alpha}}{\sqrt{2\pi} e^{-n-1} (n+1)^{n+\frac{1}{2}}} \\ &= (n+\alpha+1)^{\alpha} e^{-\alpha} \left(1 + \frac{\alpha}{n+1}\right)^{n+\frac{1}{2}} \\ &\sim (n+\alpha+1)^{\alpha} \sim n^{\alpha}. \end{aligned} \tag{68}$$

Thus

$$\begin{aligned} 1 + N \frac{\Gamma(n+\alpha+1)}{(n-1)!\Gamma(\alpha+1)\Gamma(\alpha+2)} &= 1 + \frac{\Gamma(n+\alpha+1)}{\Gamma(n)} \frac{N}{\Gamma(\alpha+1)\Gamma(\alpha+2)} \\ &\sim 1 + \frac{N n^{\alpha+1}}{\Gamma(\alpha+1)\Gamma(\alpha+2)} \\ &\sim \frac{N n^{\alpha+1}}{\Gamma(\alpha+1)\Gamma(\alpha+2)}. \end{aligned} \tag{69}$$

From (67), we have

$$\begin{aligned} u_n &= (2n + 1 + \alpha - c_n) - \frac{n(n+\alpha+1)}{c_n} \\ &= (2n + 1 + \alpha) - \frac{n + \alpha + 1 - \alpha - 1 + N \dfrac{(n+\alpha+1)\Gamma(n+\alpha+1)}{(n-1)!\Gamma(\alpha+1)\Gamma(\alpha+2)}}{1 + N \dfrac{\Gamma(n+\alpha+1)}{(n-1)!\Gamma(\alpha+1)\Gamma(\alpha+2)}} \\ &\quad -(n+\alpha+1) \frac{1 + N \dfrac{\Gamma(n+\alpha+1)}{(n-1)!\Gamma(\alpha+1)\Gamma(\alpha+2)}}{1 + N \dfrac{\Gamma(n+\alpha+2)}{n!\Gamma(\alpha+1)\Gamma(\alpha+2)}}. \end{aligned}$$

Then, after some computations, and using (68) and (69) we can estimate its behavior with respect to N and n:

$$\begin{aligned}
u_n &= \frac{\alpha+1}{\left(1+N\frac{\Gamma(n+\alpha+1)}{(n-1)!\Gamma(\alpha+1)\Gamma(\alpha+2)}\right)} - \frac{\alpha+1}{\left(1+N\frac{\Gamma(n+\alpha+2)}{n!\Gamma(\alpha+1)\Gamma(\alpha+2)}\right)} \\
&= \frac{(\alpha+1)^2}{\left(1+N\frac{\Gamma(n+\alpha+1)}{\Gamma(n+1)}\frac{n}{\Gamma(\alpha+1)\Gamma(\alpha+2)}\right)\left(1+N\frac{\Gamma(n+\alpha+1)}{\Gamma(n+1)}\frac{(n+\alpha+1)}{\Gamma(\alpha+1)\Gamma(\alpha+2)}\right)} \\
&\quad \cdot \frac{N\Gamma(n+\alpha+1)}{(n!\Gamma(\alpha+1)\Gamma(\alpha+2))} \\
&\sim \frac{(\alpha+1)^2}{\left(N\frac{n^{\alpha+1}}{\Gamma(\alpha+1)\Gamma(\alpha+2)}\right)^2} \cdot \frac{Nn^{\alpha}}{\Gamma(\alpha+1)\Gamma(\alpha+2)} \\
&= \frac{(\alpha+1)\left[\Gamma(\alpha+2)\right]^2}{N} n^{-\alpha-2}
\end{aligned}$$

The electrostatic interpretation of the distribution of zeros means that we have an equilibrium position under the presence of an external potential

$$\ln\, Q(x) + \ln\, x^{\alpha+2}e^{-x},$$

where the first term represents a short range potential corresponding to a unit charge located at u_n and the second one is a long range potential associated with the weight function (see also [10] and [11]).

2. Now, we will consider $a<0$. In this case $d\mu^*(x) = (x-a)x^{\alpha}e^{-x}dx$. Thus,

Proposition 5 *The structure relation* (44) *for the measure*

$$d\mu^*(x) = (x-a)x^{\alpha}e^{-x}dx, \quad a<0,$$

is

$$\phi(x)D(p_n^*(a;x)) = A(x,n)p_n^*(a;x) + B(x,n)p_{n-1}^*(a;x),$$

where

$$\begin{aligned}
\phi(x) &= (x-a)x, \\
A(x,n) &= n\left[x-(n+1+a_n)\left(1+\frac{n+\alpha}{a_{n-1}}\right)\right], \\
B(x,n) &= \frac{n\,(n+\alpha)}{a_{n-1}}\left[a_n x - (n+1+a_n)\,(n+1+a_n+\alpha)\right].
\end{aligned}$$

Proof From (7) we get

$$(x-a)\,p_n^*(a;x) = L_{n+1}^{\alpha}(x) - \frac{L_{n+1}^{\alpha}(a)}{L_n^{\alpha}(a)}L_n^{\alpha}(x).$$

Taking derivatives with respect to x in both hand sides of the above expression, and multiplying the resulting expression by x, we see that

$$xp_n^*(a;x) + x\,(x-a)\,D\left(p_n^*(a;x)\right) = xD\left(L_{n+1}^{\alpha}(x)\right) - \frac{L_{n+1}^{\alpha}(a)}{L_n^{\alpha}(a)}xD\left(L_n^{\alpha}(x)\right).$$

Using the structure relation (66) for Laguerre polynomials, we obtain

$$xp_n^*(a;x) + x\,(x-a)\,D\,(p_n^*(a;x)) = (n+1)\,L_{n+1}^{\alpha}(x) + (n+1)\,(n+1+\alpha)\,L_n^{\alpha}(x)$$
$$-\frac{L_{n+1}^{\alpha}(a)}{L_n^{\alpha}(a)}\left[nL_n^{\alpha}(x) + n\,(n+\alpha)\,L_{n-1}^{\alpha}(x)\right].$$

Now, from (63), we get

$$\begin{aligned}
&xp_n^*(a;x) + x\,(x-a)\,Dp_n^*(a;x)\\
&= (n+1)\,xL_n^{\alpha}(x) - (n+1)\,(2n+1+\alpha)\,L_n^{\alpha}(x) - (n+1)\,n\,(n+\alpha)\,L_{n-1}^{\alpha}(x)\\
&\quad + \left[(n+1)\,(n+1+\alpha) - \frac{nL_{n+1}^{\alpha}(a)}{L_n^{\alpha}(a)}\right] L_n^{\alpha}(x) - \frac{L_{n+1}^{\alpha}(a)}{L_n^{\alpha}(a)} n\,(n+\alpha)\,L_{n-1}^{\alpha}(x)\\
&= \left[(n+1)\,x - (n+1)\,(2n+1+\alpha) + (n+1)\,(n+1+\alpha) - \frac{nL_{n+1}^{\alpha}(a)}{L_n^{\alpha}(a)}\right] L_n^{\alpha}(x)\\
&\quad - \left(n\,(n+\alpha)\,(n+1) + n\,(n+\alpha)\,\frac{L_{n+1}^{\alpha}(a)}{L_n^{\alpha}(a)}\right) L_{n-1}^{\alpha}(x)\\
&= \left[(n+1)\,(x-n) - \frac{nL_{n+1}^{\alpha}(a)}{L_n^{\alpha}(a)}\right] L_n^{\alpha}(x)\\
&\quad - \left(n\,(n+1)\,(n+\alpha) + n\,(n+\alpha)\,\frac{L_{n+1}^{\alpha}(a)}{L_n^{\alpha}(a)}\right) L_{n-1}^{\alpha}(x).
\end{aligned}$$

Now, using the notation

$$a_n = \frac{L_{n+1}^{\alpha}\,(a)}{L_n^{\alpha}\,(a)}, \tag{70}$$

and, from (37),

$$L_n^{\alpha}(x) = p_n^*(a;x) - \frac{\gamma_n}{a_{n-1}} p_{n-1}^*(a;x),$$

we have

$$\begin{aligned}
&xp_n^*(a;x) + x\,(x-a)\,D\,(p_n^*(a;x))\\
&= ((x-n)\,(n+1) - na_n)\,L_n^{\alpha}(x) - n\,(n+\alpha)\,(n+1+a_n)\,L_{n-1}^{\alpha}(x)\\
&= ((x-n)\,(n+1) - na_n)\left[p_n^*(a;x) - \gamma_n \frac{1}{a_{n-1}} p_{n-1}^*(a;x)\right]\\
&\quad -n\,(n+\alpha)\,(n+1+a_n)\left[p_{n-1}^*(a;x) - \gamma_{n-1}\frac{1}{a_{n-2}} p_{n-2}^*(a;x)\right]\\
&= ((x-n)\,(n+1) - na_n)\,p_n^*(a;x)\\
&\quad - \left[((n+1)\,(x-n) - na_n)\,\frac{n\,(n+\alpha)}{a_{n-1}} + n\,(n+\alpha)\,(n+1+a_n)\right] p_{n-1}^*(a;x)\\
&\quad + n\,(n-1)\,(n+\alpha)\,(n-1+\alpha)\,(n+1+a_n)\,\frac{p_{n-2}^*(a;x)}{a_{n-2}}.
\end{aligned}$$

Now, using the TTRR (38) for monic kernels, and according to (39)

$$\frac{\gamma_{n-1}^{*}}{\gamma_{n-1}} = \frac{L_{n}^{\alpha}(a)}{L_{n-1}^{\alpha}(a)}\frac{L_{n-2}^{\alpha}(a)}{L_{n-1}^{\alpha}(a)}$$

we obtain

$$\begin{aligned}
\frac{a_{n-1}}{a_{n-2}} &= \frac{\gamma_{n-1}^{*}}{(n-1)\,(n-1+\alpha)} \\
\frac{(n-1)\,(n-1+\alpha)}{a_{n-2}} &= \frac{\gamma_{n-1}^{*}}{a_{n-1}}
\end{aligned}$$

and then

$$\begin{aligned}
& xp_{n}^{*}(a;x) + x\,(x-a)\,D\left(p_{n}^{*}(a;x)\right) \\
&= \left((x-n)\,(n+1) - na_{n}\right)p_{n}^{*}(a;x) \\
&\quad - \left[\left((n+1)\,(x-n) - na_{n}\right)\frac{n\,(n+\alpha)}{a_{n-1}} + n\,(n+\alpha)\,(n+1+a_{n})\right]p_{n-1}^{*}(a;x) \\
&\quad + n\,(n+\alpha)\,(n+1+a_{n})\,\frac{\gamma_{n-1}^{*}p_{n-2}^{*}(a;x)}{a_{n-1}} \\
&= \left[\left((x-n)\,(n+1) - na_{n}\right) - n\,(n+\alpha)\,(n+1+a_{n})\,\frac{1}{a_{n-1}}\right]p_{n}^{*}(a;x) \\
&\quad - n\,(n+\alpha)\left[\left((n+1)\,(x-n) - na_{n}\right)\frac{1}{a_{n-1}}\right. \\
&\quad \left. + \,(n+1+a_{n}) - (n+1+a_{n})\left(x-\beta_{n-1}^{*}\right)\frac{1}{a_{n-1}}\right]p_{n-1}^{*}(a;x) \\
&\qquad = \left[\left((n+1)\,(x-n) - na_{n}\right) - \frac{n\,(n+\alpha)\,(n+1+a_{n})}{a_{n-1}}\right]p_{n}^{*}(a;x) \\
&\qquad\quad + n\,(n+\alpha)\left[\frac{1}{a_{n-1}}\,(n+1+a_{n})\left(x-\beta_{n-1}^{*}\right)\right. \\
&\qquad\quad \left. - \frac{1}{a_{n-1}}\left((n+1)\,(x-n) - na_{n}\right) - (n+1+a_{n})\right]p_{n-1}^{*}(a;x).
\end{aligned}$$

Therefore

$$\phi(x)D(p_{n}^{*}(a;x)) = A(x,n)p_{n}^{*}(a;x) + B(x,n)p_{n-1}^{*}(a;x),$$

where

$$\begin{aligned}
\phi(x) &= x\,(x-a) \\
A(x,n) &= \left((n+1)\,(x-n) - na_{n}\right) - \frac{n\,(n+\alpha)\,(n+1+a_{n})}{a_{n-1}} - x \\
B(x,n) &= n\,(n+\alpha)\left[\frac{1}{a_{n-1}}\,(n+1+a_{n})\left(x-\beta_{n-1}^{*}\right)\right. \\
&\quad \left. - \frac{1}{a_{n-1}}\left((n+1)\,(x-n) - na_{n}\right) - (n+1+a_{n})\right].
\end{aligned}$$

Simplifying these expressions we have

$$\begin{aligned} A(x,n) &= nx - n(n+1) - na_n - \frac{n(n+\alpha)(n+1+a_n)}{a_{n-1}} \\ &= n\left[x - (n+1+a_n)\left(1+\frac{n+\alpha}{a_{n-1}}\right)\right] \end{aligned}$$

and

$$\begin{aligned} B(x.n) &= n(n+\alpha)\left[\frac{xa_n+n+n^2}{a_{n-1}} + \frac{na_n}{a_{n-1}} - \frac{(n+1+a_n)}{a_{n-1}}\beta^*_{n-1} - (n+1+a_n)\right] \\ &= n(n+\alpha)\left[\frac{a_n}{a_{n-1}}x + \frac{n+1+a_n}{a_{n-1}}\left(n-\beta^*_{n-1}\right) - (n+1+a_n)\right]. \end{aligned}$$

In the above expression, using again (39)

$$\beta^*_n = \beta_{n+1} + a_{n+1} - a_n = 2n+\alpha+3+a_{n+1}-a_n$$

we obtain

$$B(x,n) = \frac{n(n+\alpha)}{a_{n-1}}\left[a_n x - (n+1+a_n)(n+1+a_n+\alpha)\right].$$

This is an alternative approach to the method described in [16].

Notice that the Pearson equation for the linear functional associated with the measure becomes

$$D\left(\phi u^*\right) = \psi u^*$$

and (see (42) and (62))

$$\begin{aligned} \phi(x) &= (x-a)\sigma(x) = (x-a)x, \\ \psi(x) &= 2\sigma(x) + (x-a)\tau(x) = 2x + (x-a)(\alpha+1-x). \end{aligned}$$

According to (39) and (40),

$$\begin{aligned} \beta^*_{n-1} &= \beta_n + a_n - a_{n-1}, \\ \gamma^*_{n-1} &= \frac{a_{n-1}}{a_{n-2}}\gamma_{n-1}. \end{aligned}$$

This means that $Q(x)$ in (55) is the following quadratic polynomial

$$\begin{aligned} Q(x) &= B(x,n) + c_n\left[-2A(x,n) + \phi'(x) - \psi(x) + \frac{c_n}{\gamma^*_{n-1}}B(x,n-1)\right] \\ &= \gamma_n\frac{a_n}{a_{n-1}}x + \gamma_n\frac{n+1+a_n}{a_{n-1}}\left(n-(\beta_n+a_n-a_{n-1})\right) \\ &\quad -\gamma_n(n+1+a_n) - 2nc_n x + 2c_n(n+1+a_n)\left(n+\frac{\gamma_n}{a_{n-1}}\right) \\ &\quad +c_n x^2 - c_n(a+\alpha+1)x + c_n(a(\alpha+1)-a) + c_n^2 x \\ &\quad +c_n^2\frac{n+a_{n-1}}{a_{n-1}}\left(n-1-\beta_{n-1}-a_{n-1}+a_{n-2}\right) - c_n^2\frac{a_{n-2}}{a_{n-1}}(n+a_{n-1}). \end{aligned}$$

After some tedious computation the above expression becomes

$$\begin{aligned}
Q\left(x\right) &= c_n x^2 + \left(\gamma_n \frac{a_n}{a_{n-1}} - 2nc_n + c_n^2 - c_n\left(a+\alpha+1\right)\right)x \\
&\quad + \left(n+1+a_n\right)\left[\left(2c_n n - \gamma_n\right) + \frac{\gamma_n}{a_{n-1}}\left(n-\beta_n - a_n + a_{n-1} + 2c_n\right)\right] \\
&\quad + c_n^2 \frac{\left(n+a_{n-1}\right)}{a_{n-1}}\left(n-1-2n+1-\alpha-a_{n-1}\right) + ac_n\alpha,
\end{aligned}$$

i.e.,

$$Q\left(x\right) = c_n x^2 + r_n x + s_n$$

with

$$\begin{aligned}
r_n &= n\left(n+\alpha\right)\frac{a_n}{a_{n-1}} + c_n^2 - c_n\left(a+\alpha+1+2n\right) \\
&= \left(c_n+a_n\right)\left(c_n-a_n\right) - \left(c_n-a_n\right)a - \left(c_n+a_n\right)\left(2n+\alpha+1\right)
\end{aligned}$$

and

$$\begin{aligned}
s_n &= \left(n+1+a_n\right)\left[\left(n+1+a_n+\alpha\right)\left(2n+1+a_n+\alpha-a-2c_n\right)+2ac_n\right] \\
&\quad + a\alpha c_n + c_n^2\left(a_n - a_{n-1} + 1 - a\right).
\end{aligned}$$

The zeros of this polynomial are

$$\begin{aligned}
z_{1,n} &= -\frac{1}{2c_n}\left(r_n + \sqrt{r_n^2 - 4s_n c_n}\right), \\
z_{2,n} &= -\frac{1}{2c_n}\left(r_n - \sqrt{r_n^2 - 4s_n c_n}\right).
\end{aligned}$$

Taking into account

$$\begin{aligned}
\frac{\psi}{\phi} &= \frac{2\sigma(x) + (x-a)\tau(x)}{(x-a)\sigma(x)} \\
&= \frac{2}{x-a} + \frac{\tau(x)}{\sigma(x)} \\
&= \frac{2}{x-a} + \frac{\alpha+1-x}{x} \\
&= \frac{2}{x-a} + \frac{\alpha+1}{x} - 1,
\end{aligned}$$

the electrostatic interpretation means that the equilibrium position for the zeros under the presence of an external potential

$$\ln Q(x) + \ln\ (x-a)^2 x^{\alpha+1} e^{-x},$$

where the first one is a short range potential corresponding to two unit charges located at $z_{1,n}$ and $z_{2,n}$ and the second one is a long range potential associated with a polynomial perturbation of the weight function.

5.3 Electrostatic interpretation for the zeros of Jacobi type orthogonal polynomials

We shall use some basic properties of the Jacobi classical monic polynomials $P_n^{\alpha,\beta}(x)$.

i. Let u be the linear funcional

$$\langle u, p\rangle = \int_{-1}^{1} p(x)(1-x)^{\alpha}(1+x)^{\beta}dx, \quad \alpha, \beta > -1,\ p \in \mathbb{P}.$$

So u satisfies the Pearson differential equation

$$D(\sigma(x)u) = \tau(x)u,$$

where

$$\sigma(x) = 1 - x^2, \quad \tau(x) = (\beta - \alpha) - (\alpha + \beta + 2)x. \tag{71}$$

ii. For every $n \in \mathbb{N}$,

$$\begin{aligned} &P_{-1}^{\alpha,\beta}(x) = 0, \quad P_{0}^{\alpha,\beta}(x) = 1, \\ &P_{n+1}^{\alpha,\beta}(x) = (x - \beta_n)P_{n}^{\alpha,\beta}(x) - \gamma_n P_{n-1}^{\alpha,\beta}(x), \end{aligned} \tag{72}$$

where

$$\beta_n = \beta_n^{\alpha,\beta} = \frac{\beta^2 - \alpha^2}{(2n+\alpha+\beta)(2n+\alpha+\beta+2)}$$

and

$$\gamma_n = \gamma_n^{\alpha,\beta} = \frac{4n(n+\alpha)(n+\beta)(n+\alpha+\beta)}{(2n+\alpha+\beta-1)(2n+\alpha+\beta)^2(2n+\alpha+\beta+1)}.$$

iii. For every $n \in \mathbb{N}$

$$P_n^{\alpha,\beta}(-1) = \frac{(-1)^n 2^n \Gamma(n+\beta+1)\Gamma(n+\alpha+\beta+1)}{\Gamma(\beta+1)\Gamma(2n+\alpha+\beta+1)}. \tag{73}$$

iv. For every $n \in \mathbb{N}$

$$K_{n-1}(-1,-1) = \frac{1}{2^{\alpha+\beta+1}} \frac{\Gamma(n+\beta+1)\Gamma(n+\alpha+\beta+1)}{\Gamma(n)\Gamma(\beta+1)\Gamma(\beta+2)\Gamma(n+\alpha)}. \tag{74}$$

v. For every $n \in \mathbb{N}$

$$\begin{aligned} (1-x^2)D\left(P_n^{\alpha,\beta}(x)\right) &= \frac{-n[\beta - \alpha + (2n+\alpha+\beta)x]}{2n+\alpha+\beta} P_n^{\alpha,\beta}(x) \\ &\quad + \frac{4n(n+\alpha)(n+\beta)(n+\alpha+\beta)}{(2n+\alpha+\beta)^2(2n+\alpha+\beta-1)} P_{n-1}^{\alpha,\beta}(x). \end{aligned} \tag{75}$$

Now, we shall give an electrostatic interpretation for the zeros of Jacobi type polynomials $P_n^{\alpha,\beta,N}(a;x)$ which are orthogonal with respect to the measure $d\mu_N = (1-x)^{\alpha}(1+x)^{\beta}dx + N\delta_a$, that is, they are orthogonal with respect the following inner product:

$$\langle p, q\rangle = \int_{-1}^{1} p(x)q(x)(1-x)^{\alpha}(1+x)^{\beta}dx + Np(a)q(a), \quad a \notin (-1,1),\ N \geq 0.$$

We will analyze two cases:

1. Firstly, we consider $a = -1$. Thus, the polynomials $P_n^{\alpha,\beta,N}(-1;x)$ are orthogonal with respect to

$$d\mu_N = (1-x)^{\alpha}(1+x)^{\beta}dx + N\delta_{-1}.$$

Therefore, the polynomials $p_n^*(-1;x) = P_n^{\alpha,\beta+1}(x)$ associated with the measure

$$d\mu^*(x) = (x-(-1))d\mu(x) = (1-x)^{\alpha}(1+x)^{\beta+1}dx$$

have Pearson's coefficients given by (see (43) and (71))

$$\phi(x) = \sigma(x) = 1 - x^2, \quad \psi(x) = \widetilde{\sigma}(x) + \tau(x) = (\beta-\alpha+1) - (\alpha+\beta+3)\,x.$$

On the other hand, from (75), the structure relation (44) reads

$$\phi(x)D\left(P_n^{\alpha,\beta+1}(x)\right) = A(x,n)P_n^{\alpha,\beta+1}(x) + B(x,n)P_{n-1}^{\alpha,\beta+1}(x),$$

where

$$\begin{aligned} A(x,n) &= \frac{-n[\beta-\alpha+1+(2n+\alpha+\beta+1)x]}{2n+\alpha+\beta+1}, \\ B(x,n) &= \frac{4n(n+\alpha)(n+\beta+1)(n+\alpha+\beta+1)}{(2n+\alpha+\beta+1)^2(2n+\alpha+\beta)}. \end{aligned}$$

The coefficient γ_n^* in (38) when $p_n^*(-1;x) = P_n^{\alpha,\beta+1}(x)$ is

$$\gamma_n^* = \gamma_n^{\alpha,\beta+1} = \frac{4n(n+\alpha)(n+\beta+1)(n+\alpha+\beta+1)}{(2n+\alpha+\beta)(2n+\alpha+\beta+1)^2(2n+\alpha+\beta+2)}.$$

Now, we will find the coefficient c_n in (34). Using (73) and (74) it follows that

$$\begin{aligned} c_n &= -\frac{1+NK_n(-1,-1)}{1+NK_{n-1}(-1,-1)}\frac{P_{n-1}^{\alpha,\beta}(-1)}{P_n^{\alpha,\beta}(-1)} \cdot \gamma_n^{\alpha,\beta} \\ &= \frac{1+NK_n(-1,-1)}{1+NK_{n-1}(-1,-1)}\frac{2n(n+\alpha)}{(2n+\alpha+\beta)(2n+\alpha+\beta+1)} > 0, \end{aligned}$$

and, finally, we get

$$\begin{aligned} Q(x) = B(x,n) + c_n &\left[(2n+\alpha+\beta)c_n - \frac{(\alpha+\beta+1)(\beta-\alpha+1)}{2n+\alpha+\beta+1}\right] \\ &+ (2n+\alpha+\beta+1)c_n x. \end{aligned}$$

Now, we will show that the zero of $Q(x)$ belongs to $(-1,1)$. In fact, observe that

$$\begin{aligned} Q(1) &= B(x,n) + c_n\left[(2n+\alpha+\beta)c_n - \frac{(\alpha+\beta+1)(\beta-\alpha+1)}{2n+\alpha+\beta+1}\right] \\ &\quad + (2n+\alpha+\beta+1)c_n \\ &= B(x,n) + c_n\left[(2n+\alpha+\beta)c_n + \frac{2(2n(n+\alpha+\beta+1)+\alpha(\alpha+\beta+1))}{2n+\alpha+\beta+1}\right] \\ &> 0, \end{aligned}$$

and, after some tedious calculations,

$$
\begin{aligned}
Q(-1) = {} & \frac{-2^{\alpha+\beta+3}(\beta+1)\mathit{\Gamma}(n)\mathit{\Gamma}(n+\alpha)\mathit{\Gamma}(\beta+2)^2\mathit{\Gamma}(n+\beta+1)\mathit{\Gamma}(n+\alpha+\beta+1)N}{2n+\alpha+\beta} \\
& \times\frac{1}{2^{\alpha+\beta+1}\mathit{\Gamma}(n)\mathit{\Gamma}(n+\alpha)\mathit{\Gamma}(\beta+1)\mathit{\Gamma}(\beta+2)+N\mathit{\Gamma}(n+\beta+1)\mathit{\Gamma}(n+\alpha+\beta+1)} \\
& < 0.
\end{aligned}
$$

Next we will show the behavior of this zero,

$$
\begin{aligned}
u_n = & -\frac{2(n+\beta+1)(n+\alpha+\beta+1)}{(2n+\alpha+\beta+1)^2}\frac{1+NK_{n-1}(-1,-1)}{1+NK_n(-1,-1)} \\
& -\frac{2n(n+\alpha)}{(2n+\alpha+\beta+1)^2}\frac{1+NK_n(-1,-1)}{1+NK_{n-1}(-1,-1)}+\frac{(\alpha+\beta+1)(\beta-\alpha+1)}{(2n+\alpha+\beta+1)^2} \\
= & \frac{-2(n+\beta+1)(n+\alpha+\beta+1)-2n\,(n+\alpha)+(\alpha+\beta+1)(\beta-\alpha+1)}{(2n+\alpha+\beta+1)^2} \\
& +\frac{N\left(P_n^{\alpha,\beta}(-1)\right)^2/\left\|P_n^{\alpha,\beta}\right\|_\mu^2}{(2n+\alpha+\beta+1)^2}\left[\frac{2\,(n+\beta+1)\,(n+\alpha+\beta+1)}{1+NK_n\,(-1,-1)}-\frac{2n\,(n+\alpha)}{1+NK_n\,(-1,-1)}\right] \\
= & \frac{-4n^2-4\,(\alpha+\beta+1)\,n-(\alpha+\beta+1)^2}{(2n+\alpha+\beta+1)^2} \\
& +2N\frac{\left(P_n^{\alpha,\beta}(-1)\right)^2/\left\|P_n^{\alpha,\beta}\right\|_\mu^2}{(2n+\alpha+\beta+1)^2}\left[\frac{(n+\beta+1)\,(n+\alpha+\beta+1)}{1+NK_n\,(-1,-1)}-\frac{n\,(n+\alpha)}{1+NK_n\,(-1,-1)}\right] \\
& -1+2N\frac{\left(P_n^{\alpha,\beta}(-1)\right)^2/\left\|P_n^{\alpha,\beta}\right\|_\mu^2}{(2n+\alpha+\beta+1)^2}\left[\frac{(n+\beta+1)\,(n+\alpha+\beta+1)}{1+NK_n\,(-1,-1)}-\frac{n\,(n+\alpha)}{1+NK_n\,(-1,-1)}\right].
\end{aligned}
\tag{76}
$$

But from

$$
\frac{\left(P_n^{\alpha,\beta}(x)\right)^2}{\left\|P_n^{\alpha,\beta}\right\|_\mu^2}=K_n(x,x)-K_{n-1}(x,x)
$$

and (74) it easily follows that

$$
\frac{\left(P_n^{\alpha,\beta}(x)\right)^2}{\left\|P_n^{\alpha,\beta}\right\|_\mu^2}=K_{n-1}(x,x)\left(\frac{(n+\alpha+\beta+1)\,(n+\beta+1)}{n\,(n+\alpha)}-1\right). \tag{77}
$$

From (76) and (77) we get

$$\begin{aligned}
u_n &= -1+2N\frac{\left(P_n^{\alpha,\beta}(-1)\right)^2/\left\|P_n^{\alpha,\beta}\right\|_\mu^2}{(2n+\alpha+\beta+1)^2\left(1+NK_n\left(-1,-1\right)\right)}\\
&\quad\times\left[\frac{(n+\alpha+\beta+1)\,(n+\beta+1)}{n\,(n+\alpha)}-\frac{1+NK_n\left(-1,-1\right)}{1+NK_{n-1}\left(-1,-1\right)}\right]\\
&= -1+2N\frac{n(n+\alpha)\left(P_n^{\alpha,\beta}(-1)\right)^2/\left\|P_n^{\alpha,\beta}\right\|_\mu^2}{(2n+\alpha+\beta+1)^2\left(1+NK_n\left(-1,-1\right)\right)}\\
&\quad\times\left[\frac{\left(P_n^{\alpha,\beta}(-1)\right)^2/\left\|P_n^{\alpha,\beta}\right\|_\mu^2}{K_{n-1}\left(-1,-1\right)\left(1+NK_{n-1}\left(-1,-1\right)\right)}\right],\\
&> -1.
\end{aligned}$$

Moreover, using the asymptotics of $K_n\left(-1,-1\right)$, which according with (74) is

$$\begin{aligned}
&K_{n-1}(-1,-1)\\
&=\frac{1}{2^{\alpha+\beta+2}\Gamma\left(\beta+1\right)\Gamma\left(\beta+2\right)}\\
&\quad\times\frac{e^{-(n+\beta+1)}\left(n+\beta+1\right)^{\left(n+\beta+1-\frac{1}{2}\right)}e^{-(n+\alpha+\beta+1)}\left(n+\alpha+\beta+1\right)^{\left(n+\alpha+\beta+1-\frac{1}{2}\right)}}{e^{-n}n^{n-\frac{1}{2}}\left(n+\alpha\right)^{n+\alpha-\frac{1}{2}}e^{-(n+\alpha)}}\\
&\sim\frac{1}{2^{\alpha+\beta+2}\Gamma\left(\beta+1\right)\Gamma\left(\beta+2\right)}n^{2(\beta+1)},
\end{aligned}$$

and after some calculations, we finally obtain

$$\begin{aligned}
u_n &= -1+2N\frac{\left[(2\beta+2)n+(\alpha+\beta+1)\right]^2}{n(n+\alpha)\left(2n+\alpha+\beta+1\right)^2}\\
&\quad\times\frac{K_{n-1}\left(-1,-1\right)}{\left(1+NK_n\left(-1,-1\right)\right)\left(1+NK_{n-1}\left(-1,-1\right)\right)}\\
&\sim -1+\frac{2^{\alpha+\beta+2}\left(\beta+1\right)\left[\Gamma\left(\beta+2\right)\right]^2}{N}n^{-2(\beta+2)}.
\end{aligned}$$

The electrostatic interpretation means that the equilibrium position for the zeros under the presence of an external potential

$$\ln Q(x)+\ln\,(1-x)^{\alpha+1}(1+x)^{\beta+2},$$

where the first one is a short range potential corresponding to a unit charge located at the zero of $Q(x)$ and the other one is a long range potential associated with the weight function.

2. Now, we will consider $a < -1$. In this case,

$$d\mu^*(x) = (x-a)(1-x)^\alpha (1+x)^\beta dx.$$

As a consequence,

Proposition 6 *The structure relation* (44) *for the above measure is*

$$\phi(x) D(p_n^*(a;x)) = A(x,n) p_n^*(a;x) + B(x,n) p_{n-1}^*(a;x),$$

where

$$\begin{aligned}
\phi(x) &= (x-a)(1-x^2),\\
A(x,n) &= a_{n+1}(x-\beta_n) + b_{n+1} - \lambda_n a_n - (a_{n+1}\gamma_n + \lambda_n b_n)\frac{1}{\lambda_{n-1}} - 1 + x^2,\\
B(x,n) &= (a_{n+1}(x-\beta_n) + b_{n+1} - \lambda_n a_n)\,\frac{\gamma_n}{\lambda_{n-1}} + a_{n+1}\gamma_n + \lambda_n b_n\\
&\quad -(a_{n+1}\gamma_n + \lambda_n b_n)\frac{x-\beta_{n-1}^*}{\lambda_{n-1}}.
\end{aligned}$$

Proof Using the notation

$$\lambda_n = \lambda_n^{\alpha,\beta}(a) = \frac{P_{n+1}^{\alpha,\beta}(a)}{P_n^{\alpha,\beta}(a)}, \tag{78}$$

from (7) we get

$$(x-a)\, p_n^*(a;x) = P_{n+1}^{\alpha,\beta}(x) - \lambda_n P_n^{\alpha,\beta}(x).$$

Taking derivatives with respect to x in both hand sides of the above expression, and multiplying by $1-x^2$ in both members, we see that

$$\begin{aligned}
&(1-x^2) p_n^*(a;x) + (x-a)\,(1-x^2) D\,(p_n^*(a;x))\\
&= (1-x^2) D\left(P_{n+1}^{\alpha,\beta}(x)\right) - \lambda_n (1-x^2) D\left(P_n^{\alpha,\beta}(x)\right).
\end{aligned}$$

From (75),

$$(1-x^2) D\left(P_n^{\alpha,\beta}(x)\right) = a_n P_n^{\alpha,\beta}(x) + b_n P_{n-1}^{\alpha,\beta}(x),$$

where

$$a_n(x) = a_n^{\alpha,\beta}(x) = \frac{-n[\beta-\alpha+(2n+\alpha+\beta)x]}{2n+\alpha+\beta},$$

$$b_n = b_n^{\alpha,\beta} = \frac{4n(n+\alpha)(n+\beta)(n+\alpha+\beta)}{(2n+\alpha+\beta)^2(2n+\alpha+\beta-1)}.$$

Thus,

$$\begin{aligned}
&(1-x^2) p_n^*(a;x) + (x-a)\,(1-x^2) D\,(p_n^*(a;x))\\
&= a_{n+1} P_{n+1}^{\alpha,\beta}(x) + (b_{n+1} - \lambda_n a_n)\, P_n^{\alpha,\beta}(x) - \lambda_n b_n P_{n-1}^{\alpha,\beta}(x).
\end{aligned}$$

Now, from the TTRR (72) for the classical Jacobi polynomials, we get

$$\begin{aligned}
&(1-x^2) p_n^*(a;x) + (x-a)\,(1-x^2) D\,(p_n^*(a;x))\\
&= [a_{n+1}(x-\beta_n) + b_{n+1} - \lambda_n a_n]\, P_n^{\alpha,\beta}(x) - (a_{n+1}\gamma_n + \lambda_n b_n) P_{n-1}^{\alpha,\beta}(x).
\end{aligned}$$

Now, from (37) and (78),

$$P_n^{\alpha,\beta}(x) = p_n^*(a;x) - \frac{\gamma_n}{\lambda_{n-1}} p_{n-1}^*(a;x),$$

we obtain

$$\begin{aligned}
&(1-x^2)p_n^*(a;x) + (x-a)\,(1-x^2)D\,(p_n^*(a;x)) \\
&= [a_{n+1}(x-\beta_n) + b_{n+1} - \lambda_n a_n] \left[p_n^*(a;x) - \frac{\gamma_n}{\lambda_{n-1}} p_{n-1}^*(a;x)\right] \\
&\quad -(a_{n+1}\gamma_n + \lambda_n b_n) \left[p_{n-1}^*(a;x) - \frac{\gamma_{n-1}}{\lambda_{n-2}} p_{n-2}^*(a;x)\right] \\
&= [a_{n+1}(x-\beta_n) + b_{n+1} - \lambda_n a_n]\, p_n^*(a;x) \\
&\quad - \left[(a_{n+1}(x-\beta_n) + b_{n+1} - \lambda_n a_n)\,\frac{\gamma_n}{\lambda_{n-1}} + a_{n+1}\gamma_n + \lambda_n b_n\right] p_{n-1}^*(a;x) \\
&\quad (a_{n+1}\gamma_n + \lambda_n b_n)\frac{\gamma_{n-1}}{\lambda_{n-2}} p_{n-2}^*(a;x).
\end{aligned}$$

Now, using the TTRR (38) for monic kernels,

$$\begin{aligned}
&(1-x^2)p_n^*(a;x) + (x-a)\,(1-x^2)D\,(p_n^*(a;x)) \\
&= [a_{n+1}(x-\beta_n) + b_{n+1} - \lambda_n a_n]\, p_n^*(a;x) \\
&\quad - \left[(a_{n+1}(x-\beta_n) + b_{n+1} - \lambda_n a_n)\,\frac{\gamma_n}{\lambda_{n-1}} + a_{n+1}\gamma_n + \lambda_n b_n\right] p_{n-1}^*(a;x) \\
&\quad (a_{n+1}\gamma_n + \lambda_n b_n)\frac{\gamma_{n-1}}{\lambda_{n-2}} \frac{(x-\beta_{n-1}^*)p_{n-1}^*(a;x) - p_n^*(a;x)}{\gamma_{n-1}^*}.
\end{aligned}$$

According to (39) and (78), we obtain

$$\frac{\gamma_{n-1}^*}{\gamma_{n-1}} = \frac{\lambda_{n-1}}{\lambda_{n-2}}.$$

Therefore

$$\begin{aligned}
&(1-x^2)p_n^*(a;x) + (x-a)\,(1-x^2)D\,(p_n^*(a;x)) \\
&= \left[a_{n+1}(x-\beta_n) + b_{n+1} - \lambda_n a_n - (a_{n+1}\gamma_n + \lambda_n b_n)\frac{1}{\lambda_{n-1}}\right] p_n^*(a;x) \\
&\quad - \left[(a_{n+1}(x-\beta_n) + b_{n+1} - \lambda_n a_n)\,\frac{\gamma_n}{\lambda_{n-1}} + a_{n+1}\gamma_n + \lambda_n b_n \right. \\
&\qquad\qquad \left. -(a_{n+1}\gamma_n + \lambda_n b_n)\frac{x-\beta_{n-1}^*}{\lambda_{n-1}}\right] p_{n-1}^*(a;x).
\end{aligned}$$

Thus

$$\phi(x)D(p_n^*(a;x)) = A(x,n)p_n^*(a;x) + B(x,n)p_{n-1}^*(a;x),$$

where

$$\begin{aligned}
\phi(x) &= (x-a)(1-x^2),\\
A(x,n) &= a_{n+1}(x-\beta_n)+b_{n+1}-\lambda_n a_n-(a_{n+1}\gamma_n+\lambda_n b_n)\frac{1}{\lambda_{n-1}}-1+x^2,\\
B(x,n) &= (a_{n+1}(x-\beta_n)+b_{n+1}-\lambda_n a_n)\,\frac{\gamma_n}{\lambda_{n-1}}+a_{n+1}\gamma_n+\lambda_n b_n\\
&\quad -(a_{n+1}\gamma_n+\lambda_n b_n)\frac{x-\beta^*_{n-1}}{\lambda_{n-1}}.
\end{aligned}$$

Simplifying these expressions we have

$$A(x,n) = A_{n,0}+A_{n,1}x+A_{n,2}x^2$$

and

$$B(x,n) = B_{n,0}+B_{n,1}x+B_{n,2}x^2.$$

Notice that the Pearson equation for the linear functional associated with the measure

$$d\mu^*(x) = (x-a)(1-x)^\alpha(1+x)^\beta dx$$

becomes

$$D\left(\phi u^*\right) = \psi u^*,$$

with (see (42) and (71))

$$\begin{aligned}
\phi(x) &= (x-a)\sigma(x) = (x-a)(1-x^2),\\
\psi(x) &= 2\sigma(x)+(x-a)\tau(x) =\\
&\qquad 2(1-x^2)+(x-a)(\beta-\alpha-(\alpha+\beta+2)x),
\end{aligned}$$

which means that in the Jacobi case, $Q\left(x\right)$ is the following quadratic polynomial

$$\begin{aligned}
Q\left(x\right) &= B(x,n)+c_n\left[-2A(x,n)+\phi'(x)-\psi(x)+\frac{c_n}{\gamma^*_{n-1}}B(x,n-1)\right]\\
&= \left[B_{n,2}+\left(\alpha+\beta+1-2A_{n,2}+\frac{c_nB_{n-1,2}}{\gamma^*_{n-1}}\right)c_n\right]x^2\\
&\quad +\left\{\frac{c_n^2B_{n-1,1}}{\gamma^*_{n-1}}+B_{n,1}-[\alpha(a-1)+\beta(a+1)+2A_{n,1}]c_n\right\}x\\
&\quad +B_{n,0}-[2A_{n,0}+1+a(\alpha-\beta)]c_n+\frac{c_n^2B_{n-1,0}}{\gamma^*_{n-1}}.
\end{aligned}$$

Taking into account

$$\begin{aligned}
\frac{\psi}{\phi} &= \frac{2}{x-a}+\frac{\beta-\alpha-(\alpha+\beta+2)x}{1-x^2}\\
&= \frac{2}{x-a}-\frac{\alpha+1}{1-x}+\frac{\beta+1}{1+x},
\end{aligned}$$

the electrostatic interpretation means that the equilibrium position for the zeros under the presence of an external potential

$$\ln\,Q(x) + \ln\,(x-a)^2\,(1-x)^{\alpha+1}(1+x)^{\beta+1},$$

where the first one is a short range potential corresponding to two unit charges located at the zeros of $Q(x)$ and the second one is a long range potential associated with a polynomial perturbation of the weight function.

References

1. R. Álvarez-Nodarse, F. Marcellán, and J. Petronilho, *WKB Approximation and Krall-type Orthogonal Polynomials*, Acta Appl. Math. **54** (1998), 27–58.
2. C. F. Bracciali, D. K. Dimitrov, and A. Sri Ranga, *Chain sequences and symmetric generalized orthogonal polynomials*, J. Comput. Appl. Math. **143** (2002), 95–106.
3. T. S. Chihara, *An Introduction to Orthogonal Polynomials.* Mathematics and its Applications Series, Gordon and Breach, New York, 1978.
4. D. K. Dimitrov, F. Marcellán, and F. R. Rafaeli, *Monotonicity of zeros of Laguerre-Sobolev type orthogonal polynomials*, J. Math. Anal. Appl. **368** (2010), 80–89.
5. D. K. Dimitrov, M. V. Mello, and F. R. Rafaeli, *Monotonicity of zeros of Jacobi-Sobolev type orthogonal polynomials*, Appl. Numer. Math. **60** (2010), 263–276.
6. J. Dini and P. Maroni, *La multiplication d' une forme linéaire par une forme rationnelle. Application aux polynômes de Laguerre-Hahn*, Ann. Polon. Math. **52** (1990), 175–185.
7. H. Dueñas and F. Marcellán, *Laguerre-Type orthogonal polynomials. Electrostatic interpretation*, Int. J. Pure and Appl. Math. **38** (2007), 345–358.
8. H. Dueñas and F. Marcellán, *Jacobi-Type orthogonal polynomials: holonomic equation and electrostatic interpretation*, Comm. Anal. Theory Cont. Frac.**15** (2008), 4–19.
9. M. E. H. Ismail, *An electrostatics model for zeros of general orthogonal polynomials*, Pacific J. Math. **193** (2000), 355–369.
10. M. E. H. Ismail, *More on electrostatic models for zeros of orthogonal polynomials*, Numer. Funct. Anal. Optimiz. **21** (2000), 191–204.
11. M. E. H. Ismail, *Classical and Quantum Orthogonal Polynomials in One Variable*, Encyclopedia of Mathematics and its Applications, Vol. **98**. Cambridge University Press. Cambridge UK. 2005.
12. F. A. Grunbaum, *Variations on a theme of Heine and Stieltjes: an electrostatic interpretation of the zeros of certain polynomials*, J. Comput. Appl. Math. **99** (1998), 189–194.
13. R. Koekoek, *Generalizations of classical Laguerre polynomials and some q-analogues*, Doctoral Dissertation, Techn. Univ. of Delft, The Netherlands, 1990.
14. T. H. Koornwinder, *Orthogonal polynomials with weight function* $(1-x)^\alpha(1+x)^\beta + M\delta(x+1) + N\delta(x-1)$, Canad. Math. Bull. **27** (1984), 205–214.
15. F. Marcellan and P. Maroni, *Sur l' adjonction d' une masse de Dirac à une forme réguliére et semi-classique* Annal. Mat. Pura ed Appl **CLXII** (1992), 1–22.
16. F. Marcellan and A. Ronveaux, *Differential equations for classical type orthogonal polynomials*, Canad. Math. Bull. **32** (1989), 404–411.
17. P. Maroni, *Une théorie algébrique des polynômes orthogonaux. Application aux polynômes orthogonaux semi-classiques*, in *Orthogonal Polynomials and Their Applications*, C. Brezinski et al. Editors. Annals. Comput. Appl. Math. **9**. Baltzer, Basel. 1991, 95–130.
18. F. Marcellán and F. R. Rafaeli, *A Note on monotonicity of zeros of generalized Hermite-Sobolev type orthogonal polynomials.* Integral Transforms and Special Functions (2010), (doi:10.1080/10652461003714718)
19. A. F. Nikiforov and V. B. Uvarov, *Special Functions of Mathematical Physics: An unified approach*, Birkhauser Verlag, Basel. 1988.
20. J. Shohat, *A differential equation for orthogonal polynomials*, Duke Math. J. **5** (1939), 401–417.
21. G. Szegő, *Orthogonal Polynomials*, Amer. Math. Soc. Coll. Publ, Vol. **23**, 4th ed., Amer. Math. Soc., Providence, RI, 1975.

22. V. B. Uvarov, *The connection between systems of polynomials that are orthogonal with respect to different distribution functions*, J. Comput. Math. and Math. Phys. **9** (1969), 25–36.
23. A. Zhedanov, *Rational spectral transformations and orthogonal polynomials.* J. Comput. Appl. Math. **85** (1997), 67–83.
24. G. J. Yoon, *Darboux transforms and orthogonal polynomials*, Bull. Korean Math. Soc. **39** (2002), 359–376